

Original Article



Oncologic effectiveness of nerve-sparing radical hysterectomy in cervical cancer

Antonino Ditto, Giorgio Bogani , Umberto Leone Roberti Maggiore ,
Fabio Martinelli , Valentina Chiappa, Carlos Lopez, Stefania Perotto ,
Domenica Lorusso, Francesco Raspagliesi

Department of Gynecologic Oncology, IRCCS National Cancer Institute, Milan, Italy

OPEN ACCESS

Received: Aug 11, 2017

Revised: Dec 28, 2017

Accepted: Jan 31, 2018

Correspondence to

Umberto Leone Roberti Maggiore

Department of Gynecologic Oncology, IRCCS
Foundation, IRCCS National Cancer Institute,
Via Venezian 1, Milan 20133, Italy.
E-mail: umberto.leone@istitutotumori.mi.it

Copyright © 2018. Asian Society of

Gynecologic Oncology, Korean Society of
Gynecologic Oncology

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Giorgio Bogani

<https://orcid.org/0000-0001-8373-8569>

Umberto Leone Roberti Maggiore

<https://orcid.org/0000-0002-3744-2668>

Fabio Martinelli

<https://orcid.org/0000-0002-4863-1747>

Stefania Perotto

<https://orcid.org/0000-0003-0515-8299>

Conflict of Interest

No potential conflict of interest relevant to this
article was reported.

ABSTRACT

Objective: Nerve-sparing radical hysterectomy (NSRH) was introduced with the aim to reduce pelvic dysfunctions related to conventional radical hysterectomy (RH). Here, we sought to assess the effectiveness and safety of NSRH in a relatively large number of the patients of cervical cancer (CC) patients undergoing either primary surgery or neoadjuvant chemotherapy (NACT) followed by surgery.

Methods: Outcomes of consecutive patients undergoing NSRH and of a historical cohort of patients undergoing conventional RH were retrospectively reviewed.

Results: This study included 325 (49.8%) and 327 (50.2%) undergoing NSRH and RH, respectively. Via a multivariable model, nodal status was the only factor predicting for DFS (hazard ratio [HR]=2.09; 95% confidence interval [CI]=1.17–3.73; p=0.01). A trend towards high risk of recurrence was observed for patients affected by locally advanced cervical cancer (LACC) undergoing NACT followed by surgery (HR=2.57; 95% CI=0.95–6.96; p=0.06). Type of surgical procedures (NSRH vs. RH) did not influence risk of recurrence (p=0.47). Similarly, we observed that the execution of NSRH rather than RH had not a detrimental effect on OS (HR=1.19; 95% CI=0.16–9.01; p=0.87). Via multivariable model, no factor directly correlated with OS. No difference in early complication rates was observed between the study groups. Conversely, a significant higher number of late complications was reported in RH versus NSRH groups (p=0.02).

Conclusions: Our data suggested that NSRH upholds effectiveness of conventional RH, without increasing recurrence and complication rates but improving pelvic dysfunction rates.

Keywords: Uterine Cervical Neoplasms; Complications; Hysterectomy; Surgery; Survival

INTRODUCTION

Cervical cancer (CC) is currently the fourth most common malignant disease of women worldwide; according to GLOBOCAN series of the International Agency for Research on Cancer, an estimated 560,505 new cases of CC were diagnosed in 2012, and approximately 285,000 deaths due to this type of tumor were recorded during the same period [1].

Author Contributions

Conceptualization: D.A., R.F.; Data curation: B.G., L.R.M.U., M.F., C.V., L.C., P.S.; Formal analysis: B.G., L.R.M.U.; Investigation: B.G., L.R.M.U., M.F., C.V., L.C., P.S.; Methodology: B.G., L.R.M.U.; Project administration: R.F.; Supervision: D.A., L.D., R.F.; Writing - original draft: L.R.M.U.; Writing - review & editing: D.A., L.D., R.F.

Surgical treatment, including radical hysterectomy (RH) with or without bilateral salpingo-oophorectomy and pelvic lymphadenectomy +/- para-aortic lymphadenectomy, is the gold standard for women with early stage CC. Furthermore, although radio-chemotherapy represents the standard of care for patients affected by locally advanced cervical cancer (LACC), neoadjuvant chemotherapy (NACT) followed by radical surgery represents a safe alternative [2].

Ideally, surgical treatment of CC should guarantee maximal oncological disease control while minimizing early and late morbidity. Since the first description of the RH by Wertheim in 1898, several technical modifications have been developed with the aim of decreasing the rate of intra-operative complications and postoperative pelvic dysfunctions. In this scenario, the concept of preservation of autonomic nerves during RH has become the standard in many gynecological oncologic centers [3]. Nerve-sparing radical hysterectomy (NSRH) was firstly introduced and subsequently sharpened by the Japanese school [4,5]. Available studies have shown the benefits of NSRH in terms of functional outcomes (bladder, bowel, and sexual function) [6-8], while weaker evidence is available on the oncologic outcomes of this surgical procedure. In fact, a recent systematic review underlines that most of the trials lacked a control group, had small sample size, and had short follow-up intervals [9].

Therefore, the current retrospective study compared the effectiveness and safety of NSRH with conventional RH in a relatively large number of the patients of CC patients undergoing either primary surgery or NACT followed by surgery.

MATERIALS AND METHODS

A large institutional prospectively collected database, including CC patients between 1980 and 1995, and between 2001 and 2015 was retrospectively reviewed, searching records of women who underwent either primary radical surgery or NACT followed by radical surgery. The patients submitted to surgery between 1996 and 2000 were not considered in this study because in this period, according to protocol, we did not perform type III RH. All patients gave consent for the use of personal information for health research. The Institutional Review Board (IRB) of National Cancer Institute – Milan approved this study (approval number: INT-006812).

Inclusion criteria were histologically diagnosed CC, type C (both C₁ and C₂) RH according to Querleu-Morrow classification [10], and age more than 18 years.

Exclusion criteria were: other metachronous or synchronous neoplasia; history of preoperative urinary, intestinal, or sexual dysfunctions; presence of other severe comorbidities; previous pelvic surgery (all but cesarean section); presence of psychiatric disease or any condition precluding the acquisition of written informed consent.

For the purposes of the study, patients were distinguished in 2 groups: women undergoing type C₁ NSRH and those undergoing conventional type C₂ RH.

The primary outcome of this study was to evaluate the 5-year oncologic outcomes, disease-free survival (DFS) and overall survival (OS) between the study groups. Secondary outcome was to compare the rate and the type of recurrences rate. Tertiary outcome was to assess the rate of early and late complications/dysfunctions between the study groups.

Patients were preoperatively assessed with complete physical and gynecologic examination, routine blood and urine analysis, chest radiograph, and magnetic resonance imaging. Cystoscopy, rectoscopy, or computed tomography was performed only when clinically indicated. The stage of the disease was determined by the International Federation of Gynecology and Obstetrics (FIGO) staging system [11].

NACT consisted in 3 to 4 cycles of platinum-based chemotherapy. Since 2001, all patients with FIGO stage IB2, IIA2 and IIB underwent NACT. All patients included in the study underwent radical surgery including RH with or without bilateral salpingo-oophorectomy and pelvic lymphadenectomy +/- para-aortic lymphadenectomy. Type C₁ RH was performed as previously described [12-14]. Two experienced surgeons (more than 500 radical hysterectomies performed) in gynecologic oncology (F.R. and A.D.) performed all surgical procedures.

Pelvic lymphadenectomy consisted of total removal of all lymph fatty tissue around pelvic vessels and obturator fossae. Para-aortic lymphadenectomy was performed only when pelvic lymph node was positive for the presence of metastases in the iliac area on frozen section.

Pathologic evaluation included histological type, depth of cervical stromal infiltration, infiltration of parametrium and resection margins, vaginal involvement, lymph vascular space invasion (LVSI), tumor diameter, number of pelvic lymph nodes, and number of lymph node metastases.

In group RH, in case of positive lymph nodes, parametrial or/and vaginal involvement, or positive surgical resection margins (<5 mm), presence of at least 2 cervical risk factors after RH according to Sedlis criteria [15], patients underwent adjuvant radiotherapy, whereas, the presence of LVSI alone was not considered sufficient to receive adjuvant therapy. In group NSRH, in case of positive lymph nodes, parametrial or/and vaginal involvement, or positive surgical resection margins (<5 mm) patients underwent adjuvant radiotherapy. In the RH group, according to that period's protocol, all patients were submitted to adjuvant brachytherapy.

Recurrences were diagnosed during regular follow-up visits and confirmed by computed tomographic and/or magnetic resonance imaging scans. Whenever possible, histologic or cytologic confirmation was obtained. Local recurrences were defined as vaginal recurrences, pelvic recurrences, or pelvic lymph node metastases.

The following medical, perioperative and pathologic information were obtained from the database review: age, body mass index (BMI), parity, operative time, estimated blood loss, intraoperative complications, number of perioperative blood transfusions, hospital stay, early and late postoperative complications, pain evaluation using a visual analogue scale, resumption of bladder function (day of urinary catheter removal with urinary residual <100 cc after first spontaneous micturition), FIGO stage, histopathologic subtype, number and disease status of lymph nodes removed, follow-up time, DFS, OS, and locoregional recurrence (LRR). For OS, event times were computed as the interval between the date of surgery and the date of death for any cause, with censoring at the date of last follow-up for alive patients. For DFS, event times were computed as the interval between the date of surgery and the date of first relapse or death for any cause, with censoring at the date of last follow-up for event-free subjects. For LRR, event times were computed as the interval between the date of surgery and the date of local relapse as first event, censoring at the date

of last follow-up for event-free patients, and considering as competing events vaginal relapse, distant relapse, and death, whichever occurred first. Post-operative complications were arbitrarily distinguished in early (those occurring in a period of time ≤ 60 days) and late (those occurring in a period of time >60 days) complications.

1. Statistical analysis

Data are summarized using basic descriptive statistics. The normal distribution of continuous variable data was evaluated with the Kolmogorov-Smirnov test. Categorical variables were analyzed using the χ^2 test. Continuous variables, before and after treatment, were analyzed by using the paired t-test and the Wilcoxon rank sum test accordingly to data distribution.

Survival outcomes were evaluated with both Kaplan-Meier and Cox models. Hazard ratio (HR) and 95% confidence intervals (CIs) were calculated for each comparison. Univariate and multivariate analysis were performed when appropriate. All covariates with a p value less than 0.20, based on univariate analysis were included in the multivariate model. All p values were 2-sided. Statistical analysis was performed using GraphPad Prism version 6.0 for Mac (GraphPad Software, San Diego, CA, USA) and IBM-Microsoft SPSS version 20.0 for Mac (SPSS Science, Chicago, IL, USA). The p values <0.05 were statistically significant.

RESULTS

The study included 652 patients: 325 (49.8%) and 327 (50.2%) undergoing NSRH and RH, respectively. The median (range) follow-up was 38 months (range, 1–131 months) and 123 months (range, 1–338 months) for NSRH and RH group, respectively. Main characteristics of the 2 study groups are shown in **Table 1**. No significant difference in the histology was reported between the 2 study groups. Most patients had squamous cell carcinoma, 74.5% and 87.8% for NSRH and RH, respectively, while for adenocarcinoma, these figures were 15.7% and 7.4%. The pathological analysis showed that the rate of positive pelvic nodes was similar between NSRH (20.9%) and RH (19.9%) groups. No significant difference in the median (range) number of pelvic/para-aortic lymph nodes dissected was reported between NSRH (37.5 [12–80]) and RH (36 [10–105]) groups ($p=0.546$). The large majority of patients included in the NSRH group (89%) received NACT in comparison with RH group (21.1%; $p<0.01$).

1. Recurrences and survival outcomes

Considering all the patients, 5-year DFS (**Fig. 1A**) was significantly inferior in NSRH group (77.7%) compared with RH group (84.5%; log-rank=5.394; $p=0.02$); 5-year OS (**Fig. 1B**) was 90.4% in NSRH group and 84.9% in RH group (log-rank=2.214; $p=0.13$). Interestingly, considering only patients who received NACT, 5-year DFS (**Fig. 1C**) in NSRH and RH groups were 76.6% and 75.6% (log rank=0.000; $p=0.98$), respectively; 5-year OS (**Fig. 1D**) was 89.3% in NSRH group and 79.4% in RH group (log rank=3.664; $p=0.06$).

Via a multivariable model (**Table 2**), it was observed that nodal status is the only factor predicting for DFS (HR=2.09; 95% CI=1.17–3.73; $p=0.01$). Type of surgical procedures (NSRH vs. RH) did not influence risk of recurrence ($p=0.47$). The execution of NSRH did not correlate with an increased risk of developing local (HR=1.32; 95% CI=0.58–3.01; $p=0.50$), regional (HR=1.26; 95% CI=0.41–3.81; $p=0.67$) and distant (HR=0.34; 95% CI=0.08–1.35; $p=0.12$) recurrences. Via multivariable model (**Table 3**), no factor directly correlated with OS. Furthermore, we observed that the execution of NSRH rather than RH had not a

Radical hysterectomy in cervical cancer

Table 1. Main characteristics of the study population

Characteristics	Group NSRH (n=325)	Group RH (n=327)	p
Age (yr)	49 (26–77)	46 (22–75)	
BMI			0.85
Normal (< 30)	258 (79.4)	256 (78.3)	
Obese (≥30 and <35)	24 (7.4)	28 (8.5)	
Severely obese (≥35)	12 (3.7)	15 (4.6)	
Missing data	31 (9.5)	28 (8.6)	
Histology			<0.01
Squamous cell	242 (74.5)	287 (87.8)	
Adenocarcinoma	51 (15.7)	24 (7.4)	
Other type	19 (5.8)	4 (1.2)	
Missing data	13 (4.0)	12 (3.6)	
Grade			<0.01
1	14 (4.3)	8 (2.5)	
2	98 (30.2)	23 (7.0)	
3	187 (57.5)	23 (7.0)	
Missing data	26 (8.0)	273 (83.5)	
FIGO stage			<0.01
IA2	1 (0.3)	2 (0.6)	
IB1	51 (15.7)	173 (52.9)	
IB2	88 (27.1)	83 (25.4)	
IIA	57 (17.5)	44 (13.5)	
IIB	128 (39.4)	25 (7.6)	
LVSI			<0.01
No	134 (41.2)	6 (1.8)	
Yes	178 (54.8)	53 (16.2)	
Missing data	13 (4.0)	268 (82.0)	
Parametrium			<0.01
Negative	279 (85.8)	302 (92.4)	
Positive	46 (14.2)	25 (7.6)	
Vagina			<0.01
Negative	254 (78.2)	289 (88.4)	
Positive	71 (21.8)	38 (11.6)	
Lymph node			0.74
Negative	257 (79.1)	262 (80.1)	
Positive	68 (20.9)	65 (19.9)	
Pelvic lymphadenectomy			1.00
Performed	324 (99.7)	325 (99.4)	
Sampling	1 (0.3)	2 (0.6)	
Not performed	0 (0)	0 (0)	
Para-aortic lymphadenectomy			<0.01
Performed	31 (9.6)	94 (28.8)	
Sampling	17 (5.2)	2 (0.6)	
Not performed	277 (85.2)	231 (70.6)	
Salpingo-oophorectomy			0.67
No	112 (34.5)	91 (27.8)	
Yes	213 (65.5)	236 (72.2)	
Depth of stromal invasion in the specimen (mm)			<0.01
<3	111 (34.2)	62 (19.0)	
3–10	106 (32.6)	151 (46.2)	
>10	108 (33.2)	114 (34.8)	
NACT			<0.01
No	28 (8.6)	258 (78.9)	
Yes	295 (89.0)	69 (21.1)	
Post-surgical therapy			<0.01
None	167 (51.4)	31 (9.5)	
Radiotherapy	83 (25.5)	291 (89.0)	
Chemotherapy	24 (7.4)	4 (1.2)	
Chemoradiotherapy	51 (15.7)	1 (0.3)	

Data are presented as median (range) or number (percentage).

BMI, body mass index; FIGO, International Federation of Gynecology and Obstetrics; LVSI, lymph vascular space invasion; NACT, neoadjuvant chemotherapy; NS, not significant; NSRH, nerve-sparing radical hysterectomy; RH, radical hysterectomy.




Fig. 1. Five-year DFS and OS in the study population and in the NACT subgroup of patients (time is expressed in months). (A) 5-year DFS in NSRH and RH groups. (B) 5-year OS in NSRH and in RH groups. (C) 5-year DFS considering only patients who received NACT in NSRH and RH groups. (D) 5-year OS considering only patients who received NACT in NSRH and RH groups.
DFS, disease-free survival; NACT, neoadjuvant chemotherapy; NSRH, nerve-sparing radical hysterectomy; OS, overall survival; RH, radical hysterectomy.

detrimental effect on OS (HR=1.19; 95% CI=0.16–9.01; p=0.87). Furthermore, no difference either in the rate or in the type of recurrence was reported between the two study groups (**Supplementary Table 1**).

2. Surgery related outcomes

A detailed presentation of surgical characteristics is described in **Table 4**. No statistical difference between the two study groups was observed in the operative time. Blood loss as well the number of blood units transfused was superior in patients of RH group than those of NSRH group. Median (range) hospital stay was shorter for women of NSRH group (7 [3-16]) compared with those of RH group (18 [3-131]; p<0.01). Higher rates of intraoperative complications were reported in RH group (n=65) versus NSRH group (n=41; p=0.01). No difference in early complication rates was observed between the 2 study groups. Conversely,

Radical hysterectomy in cervical cancer

Table 2. Factors influencing DFS

Characteristics	Univariate analysis for DFS	p	Multivariate analysis for DFS	p
Age (yr)	0.99 (0.97–1.01)	0.35	-	-
BMI (kg/m ²)	1.04 (0.69–1.55)	0.83	-	-
Parity	0.96 (0.76–1.21)	0.74	-	-
Histology		0.37		-
Squamous cell	Reference			
Non-squamous cell	0.62 (0.21–1.77)		-	
FIGO grade		<0.01		0.25
Grade 1&2	Reference		Reference	
Grade 3	1.97 (1.27–3.07)		1.36 (0.81–2.28)	
LVSI		0.02		0.50
No	Reference		Reference	
Yes	1.83 (1.09–3.09)		1.23 (0.67–2.23)	
FIGO stage at presentation		<0.01		0.06
Early stage	Reference		Reference	
Locally advance stage	1.78 (1.20–2.64)		2.57 (0.95–6.96)	
Nerve sparing		0.02		0.47
No	Reference		Reference	
Yes	1.53 (1.05–2.22)		0.67 (0.23–2.00)	
Parametrial involvement		<0.01		0.25
No	Reference		Reference	
Yes	1.53 (1.05–4.03)		1.46 (0.77–2.80)	
Vaginal involvement		<0.01		-
No	Reference		Reference	
Yes	2.83 (1.91–4.20)		1.30 (0.73–2.33)	
Lymph nodes positivity		<0.01		0.01
No	Reference		Reference	
Yes	3.38 (2.35–4.88)		2.09 (1.17–3.73)	

Data are presented as HR (95% CI).

BMI, body mass index; DFS, disease-free survival; FIGO, International Federation of Gynecology and Obstetrics; LVSI, lymph vascular space invasion.

Table 3. Factors influencing OS

Characteristics	Univariate analysis for OS	p	Multivariate analysis for OS	p
Age (yr)	1.01 (0.99–1.02)	0.36	-	-
BMI (kg/m ²)	1.03 (0.79–1.25)	0.76	-	-
Parity	0.95 (0.89–1.11)	0.89	-	-
Histology		0.01		0.86
Squamous cell	Reference		Reference	
Non-squamous cell	0.33 (0.14–7.60)		0.27 (0.63–1.20)	
FIGO grade		<0.01		0.88
Grade 1&2	Reference		Reference	
Grade 3	2.86 (1.38–5.97)		2.53 (0.87–7.34)	
LVSI		<0.01		0.80
No	Reference		Reference	
Yes	5.03 (1.78–14.1)		2.77 (0.89–8.64)	
FIGO stage at presentation		0.68		-
Early stage	Reference			
Locally advance stage	1.09 (0.72–1.64)		-	
Nerve sparing		0.06		0.87
No	Reference		Reference	
Yes	0.66 (0.40–1.07)		1.19 (0.16–9.01)	
Parametrial involvement		0.02		0.79
No	Reference		Reference	
Yes	1.92 (1.07–3.47)		1.14 (0.42–3.09)	
Vaginal involvement		<0.01		0.49
No	Reference		Reference	
Yes	2.18 (1.35–3.53)		1.37 (0.57–3.29)	
Lymph nodes positivity		<0.01		0.10
No	Reference		Reference	
Yes	3.05 (1.35–3.53)		2.05 (0.87–4.84)	

Data are presented as HR (95% CI).

BMI, body mass index; FIGO, International Federation of Gynecology and Obstetrics; LVSI, lymph vascular space invasion; OS, overall survival.

Table 4. Intraoperative and postoperative characteristics

Characteristics	Group NSRH (n=325)	Group RH (n=327)	p
Operative time (min)	240 (138–370)	240 (120–365)	0.42
Blood loss (mL)	300 (50–2,500)	600 (100–3,000)	<0.01
Number of blood units transfused	1 (0–5)	2 (0–10)	<0.01
Hospital stay (day)	7 (3–16)	18 (7–131)	<0.01
Number of intraoperative complications*	5 (1.5)	11 (3.4)	0.13
Bladder function			<0.01
No recovery	16 (4.9)	43 (13.1)	
Recovery	309 (95.1)	275 (84.1)	
Missing data	0 (0)	9 (2.8)	
Presence of postoperative complications (grade ≥3) [†]	41 (12.6)	65 (19.8)	0.01
Total early postoperative complications (grade ≥3)	21 (6.5)	28 (8.6)	0.40
Early postoperative complications (grade ≥3)			0.04
Lymphatic (lymphocyst-lymphorrhea)	3 (0.9)	13 (4.0)	
Hemorrhage/bleeding	9 (2.8)	2 (0.6)	
Intestinal ileus	1 (0.3)	3 (0.9)	
Urological	2 (0.6)	3 (0.9)	
Thromboembolism	0 (0)	2 (0.6)	
Need for another surgery	1 (0.3)	2 (0.6)	
Infection	1 (0.3)	2 (0.6)	
Neurologic	3 (0.9)	1 (0.3)	
Voiding	1 (0.3)	0 (0)	
More than one complication	0 (0)	0 (0)	
Total late postoperative complications (grade ≥3)	20 (6.2)	37 (11.3)	0.02
Late postoperative complications (grade ≥3)			0.05
Lymphatic (lymphocyst-lymphorrhea)	6 (1.8)	12 (3.7)	
Impaired bowel function	0 (0)	9 (2.8)	
Urological	7 (2.2)	6 (1.8)	
Need for another surgery	4 (1.2)	4 (1.2)	
Infection	1 (0.3)	4 (1.2)	
Voiding	1 (0.3)	2 (0.6)	
More than one complication	1 (0.3)	0 (0)	

Data are presented as median (range) or number (percentage).

*According to the Common Terminology Criteria for Adverse Events (CTCAE); [†]some patients had both early and late postoperative complications.

a significant higher number of late complications was reported in RH group (n=37) versus NSRH group (n=20; p=0.02).

DISCUSSION

The current study investigated the effectiveness and safety of NSRH with RH in a relatively large number of the patients of CC patients undergoing either primary surgery or NACT followed by surgery. It was reported no difference in 5-year OS was significantly between RH and NSRH groups, while 5-year DFS was superior in RH group compared with NSRH group. However, this finding is not surprising due to the higher number of patients of NSRH group with LACC and treated with NACT, and to the larger use of adjuvant radiotherapy in NSRH group. In fact, when considering only patients receiving NACT no difference in both 5-year DFS and OS was reported. Interestingly, while no difference has been reported in the rate of early complications between 2 study groups, a higher number of late complications was experienced by patients of RH group, mainly due to lymphatic complications or intestinal ileus. Patients belonging to this group were operated in an older period of time compared with patients of NSRH group; therefore, the use of vascular clips and energy coagulation was superior in NSRH group and this may justify the higher number of lymphatic complications in RH group. In addition, it is well known that the autonomic nerves innervate the bowel and

that the damage to these nerves in the pelvis can lead to diarrhoea or constipation [12,16,17], thus explaining why more patients experienced impaired bowel function in RH group.

Different trials have compared class II RH with class III NSRH showing similar results in terms of bladder function, quality of life, and oncological outcomes. However, only few studies have investigated the outcomes of type C₁ NSRH compared with conventional type C₂ RH in the treatment of CC [14,18]. Our previous research included a total of 496 patients, of whom 185 patients were treated with type C₁ NSRH and 311 patients with type C₂ RH with a median follow-up of 93 months (42 and 159 months for the NSRH and RH groups, respectively). This study demonstrated that the oncologic results and relapse rates were comparable between the 2 study groups. Bladder function and postoperative complications rate were significantly better in the NSRH than in RH group [14]. These findings are confirmed by the current study, which was conducted in a substantially enlarged study population, supporting the role of nerve-sparing technique. More recently a randomized controlled trial (RCT) was conducted to evaluate the efficacy of type C₁ NSRH in preserving bladder function and its oncologic safety in the treatment of CC. Ninety-two patients with CC stage IA2 to IIA were randomized to surgical treatment with type C₂ RH (N=44) or type C₁ NSRH (n=48), and 86 patients finally included in the analysis. This trial demonstrated no difference in 10-year DFS between the RH group (94.9%) and NSRH group (92.4%). Furthermore, volume of residual urine and bladder compliance were significantly deteriorated at 12 months after RH, while all parameters of urodynamic study were recovered no later than 3 months after NSRH. This study is the only available RCT investigating the issue of type C₁ NSRH versus conventional type C₂ RH. However, from a merely statistical point of view, it is significantly underpowered to assess oncologic outcomes. Indeed, a recent systematic review on this issue by Basaran and colleagues [9] has been underlined this concept. In their paper, the Authors calculated that the number of patients needed to prove non-inferiority of NSRH to RH with regard to recurrence would be between 4,300 and 1,000 depending on the expected risk of recurrence (5% and 20%, respectively) [9]. In this view, we deem that our study provides important data due to the large number of patients included; however, we acknowledge that the main limitations of the current work are represented by the inherent biases of the retrospective, single centre study design. Another limitation of this study is that standardized questionnaires and/or urodynamic study were not used to evaluate bladder function before and after surgery, thus limiting the assessment of this specific outcome. Finally, the significant difference in length of follow-up between the two study groups prevents us to show the long-term follow-up (10 years) of the patients.

In conclusion, this study supports the efficacy and safety of NSRH for the treatment of CC patients. NSRH seems superior to RH, NSRH correlates with lower morbidity rate and similar oncologic outcomes than RH. Although the retrospective design of the study, we deem that our findings may be useful to provide a more detailed preoperative and postoperative counselling to the patients. However, well-designed, large-scale prospective non-inferiority trials are strongly required to confirm these results.

SUPPLEMENTARY MATERIAL

Supplemental Table 1

Rate and type of recurrences

[Click here to view](#)

REFERENCES

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-86.
[PUBMED](#) | [CROSSREF](#)
2. Raspagliesi F, Bogani G, Martinelli F, Signorelli M, Chiappa V, Scaffa C, et al. Incorporating 3D laparoscopy for the management of locally advanced cervical cancer: a comparison with open surgery. *Tumori* 2016;102:393-7.
[PUBMED](#) | [CROSSREF](#)
3. Rob L, Halaska M, Robova H. Nerve-sparing and individually tailored surgery for cervical cancer. *Lancet Oncol* 2010;11:292-301.
[PUBMED](#) | [CROSSREF](#)
4. Kobayashi T. Abdominal radical hysterectomy with pelvic lymphadenectomy for cancer of the cervix. Tokyo: Nanzando; 1961.
5. Fujii S. Anatomic identification of nerve-sparing radical hysterectomy: a step-by-step procedure. *Gynecol Oncol* 2008;111:S33-41.
[PUBMED](#) | [CROSSREF](#)
6. Espino-Strebel EE, Luna JT, Domingo EJ. A comparison of the feasibility and safety of nerve-sparing radical hysterectomy with the conventional radical hysterectomy. *Int J Gynecol Cancer* 2010;20:1274-83.
[PUBMED](#) | [CROSSREF](#)
7. Cibula D, Velechovska P, Sláma J, Fischerova D, Pinkavova I, Pavlista D, et al. Late morbidity following nerve-sparing radical hysterectomy. *Gynecol Oncol* 2010;116:506-11.
[PUBMED](#) | [CROSSREF](#)
8. Bogani G, Serati M, Nappi R, Cromi A, di Naro E, Ghezzi F. Nerve-sparing approach reduces sexual dysfunction in patients undergoing laparoscopic radical hysterectomy. *J Sex Med* 2014;11:3012-20.
[PUBMED](#) | [CROSSREF](#)
9. Basaran D, Dusek L, Majek O, Cibula D. Oncological outcomes of nerve-sparing radical hysterectomy for cervical cancer: a systematic review. *Ann Surg Oncol* 2015;22:3033-40.
[PUBMED](#) | [CROSSREF](#)
10. Querleu D, Morrow CP. Classification of radical hysterectomy. *Lancet Oncol* 2008;9:297-303.
[PUBMED](#) | [CROSSREF](#)
11. FIGO Committee on Gynecologic Oncology. FIGO staging for carcinoma of the vulva, cervix, and corpus uteri. *Int J Gynaecol Obstet* 2014;125:97-8.
[PUBMED](#) | [CROSSREF](#)
12. Raspagliesi F, Ditto A, Kusamura S, Fontanelli R, Spatti G, Solima E, et al. Nerve-sparing radical hysterectomy: a pilot study. *Tumori* 2003;89:497-501.
[PUBMED](#)
13. Raspagliesi F, Ditto A, Fontanelli R, Solima E, Hanozet F, Zanaboni F, et al. Nerve-sparing radical hysterectomy: a surgical technique for preserving the autonomic hypogastric nerve. *Gynecol Oncol* 2004;93:307-14.
[PUBMED](#) | [CROSSREF](#)
14. Ditto A, Martinelli F, Mattana F, Reato C, Solima E, Carcangiu M, et al. Class III nerve-sparing radical hysterectomy versus standard class III radical hysterectomy: an observational study. *Ann Surg Oncol* 2011;18:3469-78.
[PUBMED](#) | [CROSSREF](#)
15. Sedlis A, Bundy BN, Rotman MZ, Lentz SS, Mudderspach LI, Zaino RJ. A randomized trial of pelvic radiation therapy versus no further therapy in selected patients with stage IB carcinoma of the cervix after radical hysterectomy and pelvic lymphadenectomy: a Gynecologic Oncology Group Study. *Gynecol Oncol* 1999;73:177-83.
[PUBMED](#) | [CROSSREF](#)
16. Pieterse QD, Maas CP, ter Kuile MM, Lowik M, van Eijkeren MA, Trimbos JB, et al. An observational longitudinal study to evaluate miction, defecation, and sexual function after radical hysterectomy with pelvic lymphadenectomy for early-stage cervical cancer. *Int J Gynecol Cancer* 2006;16:1119-29.
[PUBMED](#) | [CROSSREF](#)
17. van Gent MD, Romijn LM, van Santen KE, Trimbos JB, de Kroon CD. Nerve-sparing radical hysterectomy versus conventional radical hysterectomy in early-stage cervical cancer. A systematic review and meta-analysis of survival and quality of life. *Maturitas* 2016;94:30-8.
[PUBMED](#) | [CROSSREF](#)
18. Roh JW, Lee DO, Suh DH, Lim MC, Seo SS, Chung J, et al. Efficacy and oncologic safety of nerve-sparing radical hysterectomy for cervical cancer: a randomized controlled trial. *J Gynecol Oncol* 2015;26:90-9.
[PUBMED](#) | [CROSSREF](#)