

INFINITESIMAL NEWTON–OKOUNKOV BODIES AND JET SEPARATION

ALEX KÜRONYA AND VICTOR LOZOVANU

INTRODUCTION

In this paper we wish to continue the investigations initiated in [KL14, KL15] to find a satisfactory theory of positivity for divisors in terms of convex geometry. To be more specific, our aim here is to relate local positivity of line bundles to Newton–Okounkov bodies attached to infinitesimal flags.

Ever since the advent of Newton–Okounkov bodies in projective geometry (see [KKh, LM] or the review [B] for an introduction) the main question has been how their geometry is connected to the properties of the underlying polarized variety. For example, attention has been devoted to the combinatorial study of Newton–Okounkov bodies in terms of projective data (see for instance [AKL, KLM, LSS, PSU]). Nevertheless, in order for these invariants to be really useful in the quest for understanding projective varieties, it is more important to uncover implications in the other direction, that is, one should be able to gain information about line bundles in terms of their Newton–Okounkov bodies.

The hope for such results comes from Jow’s theorem [J] claiming that the function associating to an admissible flag the Newton–Okounkov body of a given divisor determines the latter up to numerical equivalence. Following our earlier work [KL14, KL15], we are interested in a local version of Jow’s principle: we will be mostly concerned with the situation where all flags considered are centered at a given point of the variety.

Compared to [KL15] we specialize the flags further; as suggested by [KL14, Sections 3 & 4], one can obtain particularly precise results by taking linear flags in the exceptional divisor of the blow-up of the point. This way, we can not only achieve a description of ampleness and nefness in terms of infinitesimal Newton–Okounkov bodies, but are also able to extend the convex geometric interpretation of moving Seshadri constants described in [KL14] to all dimensions.

Let now X be a projective variety over the complex numbers, L a big line bundle, and $x \in X$ a closed point. We say that L is locally positive or locally ample at x if there exists a neighbourhood $\mathcal{U} \subseteq X$ such that the Kodaira map ϕ_{mL} restricted to \mathcal{U} is an embedding for all $m \gg 0$. One can of course work with the alternative description provided by global generation of large twists of coherent sheaves (cf. [PAG1, Example 1.2.21] and [K, Proposition 2.7]), in any case both conditions end up being equivalent to x belonging to the complement of the augmented base locus $\mathbf{B}_+(L)$ of L (see [BCL, Theorem A]).

Once a line bundle L has been proven to be locally positive at a point $x \in X$, one can try to measure the extent of its positivity there. The traditional way to do this is via the Seshadri constant $\varepsilon(L; x)$ introduced by Demailly [D] (see also [PAG1, Chapter 5] for a thorough introduction and an extensive bibliography), or, in our setting, its extension, the moving Seshadri constant $\varepsilon(\|L\|; x)$ developed by Nakamaye [N], and studied in much more detail by Ein–Lazarsfeld–Mustață–Nakamaye–Popa [ELMNP2].

Since one can describe both local ampleness and moving Seshadri constants in terms of infinitesimal Newton–Okounkov bodies, the convex-geometric picture of local positivity appears to be complete. The first main result of our work is a characterization of ampleness and nefness in terms of Newton–Okounkov bodies (cf. [KL14, Theorem A] and [KL15, Theorems A & B], see also [CHPW]).

To fix terminology, let X be a smooth projective variety of dimension n , $x \in X$ a closed point, and $\pi: X' \rightarrow X$ be the blow-up of X at x with exceptional divisor E . An infinitesimal flag Y_\bullet over x is an admissible flag

$$Y_\bullet : Y_0 = X' \supseteq Y_1 = E \supseteq Y_2 \supseteq \dots \supseteq Y_n,$$

where each Y_i is a linear subspace of $E \simeq \mathbb{P}^{n-1}$ of dimension $n-i$ for each $i = 2, \dots, n$. The Newton–Okounkov body of π^*D with respect to Y_\bullet on X' will be denoted by $\tilde{\Delta}_{Y_\bullet}(D)$. For further results regarding infinitesimal Newton–Okounkov bodies the reader is kindly referred to Section 2.

Theorem A. (Corollary 3.3) Let X be a smooth projective variety of dimension n , D a big \mathbb{R} -divisor on X . Then the following are equivalent.

- (1) D is nef.
- (2) For every point $x \in X$ there exists an infinitesimal flag Y_\bullet over x such that $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D)$.
- (3) One has $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D)$ for every infinitesimal flag over X .

Before we proceed, let us define what we call the inverted standard simplex of size $\xi > 0$: this is the convex body

$$\Delta_\xi^{-1} \stackrel{\text{def}}{=} \text{convex hull of } \{\mathbf{0}, \xi \mathbf{e}_1, \xi(\mathbf{e}_1 + \mathbf{e}_2), \dots, \xi(\mathbf{e}_1 + \mathbf{e}_n)\} \subseteq \mathbb{R}^n,$$

where $\mathbf{e}_1, \dots, \mathbf{e}_n$ denote the standard basis vectors for \mathbb{R}^n . Lemma 2.4 and Proposition 2.6 below explain how the polytopes Δ_ξ^{-1} arise very naturally in the infinitesimal setting.

Theorem B. (Corollary 4.2) Let X be a smooth projective variety of dimension n , D a big \mathbb{R} -divisor on X . Then the following are equivalent.

- (1) D is ample.
- (2) For every point $x \in X$ there exists an infinitesimal flag Y_\bullet over x and a real number $\xi > 0$ for which $\Delta_\xi^{-1} \subseteq \tilde{\Delta}_{Y_\bullet}(D)$.
- (3) $\tilde{\Delta}_{Y_\bullet}(D)$ contains a non-trivial inverted standard simplex for every infinitesimal flag Y_\bullet over X .

Note that as opposed to [KL15, Theorem B], the theorem above provides a full generalization of what happens in the surface case; its proof is significantly more difficult than that of any of its predecessors.

An interesting feature of the argument leading to Theorem B is that it passes through separation of jets. In fact, an important step in the proof is Proposition 4.9 which claims that line bundles whose infinitesimal Newton–Okounkov bodies contain large inverted standard simplices will separate many jets. Not surprisingly, we will make an extensive use of the circle of ideas around jet separation and moving Seshadri constants, and with it, the non-trivial results of [ELMNP2]. Another important ingredient of the proof is an acute observation of Fulger–Kollár–Lehmann [FKL, Theorem A] linking inequalities between volumes of divisors to augmented base loci.

It follows from our argument that infinitesimal Newton–Okounkov bodies on projective varieties always contain inverted standard simplices at points where the divisor is locally ample. Given an infinitesimal flag Y_\bullet , the supremum of the sizes of all such is called the inverted largest simplex constant, and will be denoted by $\xi_{Y_\bullet}(D; x)$. It will turn out that this constant does not depend on the choice of the infinitesimal flag taken, leading to the common value $\xi(D; x)$. As a result of our efforts we obtain a description of moving Seshadri constants in all dimensions (cf. [KL14, Theorem D]) in the following form.

Theorem C. Let D be a big \mathbb{R} -divisor on a smooth projective variety X , $x \notin \mathbf{B}_+(D)$. Then

$$\varepsilon(\|D\|; x) = \xi(D; x) .$$

Beside providing an alternative way of defining moving Seshadri constants, the largest inverted simplex constant has other benefits as well. Via Theorem 3.1 and Theorem 4.1 it explains quite clearly why $\varepsilon(\|D\|; x) = 0$ for a divisor D with $x \in \mathbf{B}_+(D) \setminus \mathbf{B}_-(D)$.

An interesting by-product of our result is a statement about the existence of global sections with prescribed vanishing behaviour. From the definition of Newton–Okounkov bodies it is a priori quite unclear which rational points arise as actual images of global sections, and in general it is very difficult to decide when it comes to boundary points. As it turns out, for infinitesimal Newton–Okounkov bodies the situation is more amenable.

Corollary D. (Corollary 4.13) Let D be a big \mathbb{Q} -divisor on X , $x \in X$ a closed point, and Y_\bullet an infinitesimal flag over x . If $\Delta_\xi^{-1} \subseteq \Delta_{Y_\bullet}(\pi^*(D))$ for some $\xi > 0$, then all vectors in $\Delta_\xi^{-1} \cap \mathbb{Q}^n$ not lying on the face generated by the points $\lambda \cdot \mathbf{e}_1, \lambda(\mathbf{e}_1 + \mathbf{e}_2), \dots, \lambda(\mathbf{e}_1 + \mathbf{e}_n)$ are valutive.

Finally, a somewhat tentative side remark regarding moving Seshadri constants and asymptotic multiplicities. For a given point $x \in X$, the loci of \mathbb{R} -divisor classes in $\text{Big}(X)$ where $\varepsilon(\|D\|; x)$ and $\text{mult}_x \|D\|$ are naturally defined are complementary, and we point out that one can glue these functions to a unique one via

$$\varepsilon_x(D) \stackrel{\text{def}}{=} \begin{cases} \varepsilon(\|D\|; x) & \text{if } x \notin \mathbf{B}_+(D) \\ 0 & \text{if } x \in \mathbf{B}_+(D) \setminus \mathbf{B}_-(D) \\ -\text{mult}_x \|D\| & \text{if } x \in \mathbf{B}_-(D) , \end{cases}$$

which ends up being homogeneous of degree one and continuous on the big cone, while examples suggest that one can hope for ε_x to be concave. We believe that ε_x could prove useful as an extension of the moving Seshadri constant function by being capable of distinguishing between divisor classes D with $x \in \mathbf{B}_+(D) \setminus \mathbf{B}_-(D)$ and $x \in \mathbf{B}_-(D)$. In the end we discuss an example where the Seshadri function is not everywhere differentiable on the ample cone.

A few words about the organization of the paper. We begin in Section 1 by fixing notation and collecting useful facts about asymptotic base loci, Newton–Okounkov bodies, and moving Seshadri constants, in Section 2 we present some important observations about infinitesimal Newton–Okounkov bodies. The characterization of restricted base loci is given in Section 3, while Section 4 is devoted to the main part of the paper, the description of augmented base loci in terms of Newton–Okounkov bodies with the help of separation of jets. Lastly, Section 5 hosts the discussion on Seshadri functions.

Acknowledgements. We are grateful to Mihnea Popa for helpful discussions, and to the Deutsche Bahn, the Österreichische Bundesbahn, the SNCF and Thello for providing us with excellent working conditions.

1. NOTATION AND PRELIMINARIES

1.1. Notation. We work over the complex number field, X will stand for a projective variety of dimension n which will often be taken to be smooth. The point $x \in X$ will always be assumed a smooth point, while all points on varieties are taken to be closed. A divisor is always Cartier, whether it is integral, \mathbb{Q} -, or \mathbb{R} -Cartier and D will denote a big divisor without exception.

If F is an effective \mathbb{R} -Cartier divisor on X , then we write

$$\mu_F(D) = \mu(D; F) \stackrel{\text{def}}{=} \sup \{ t > 0 \mid D - tF \text{ is big} \} .$$

Furthermore, if $Z \subseteq X$ is a smooth subvariety, then denote by

$$\mu_Z(D) = \mu(D; Z) \stackrel{\text{def}}{=} \mu(\pi^* D; E) ,$$

where $\pi: X' \rightarrow X$ denotes the blow-up of X along Z with exceptional divisor E .

Remark 1.1. Based on the definition of moving Seshadri constant given below, it is not hard to see that $0 < \varepsilon(\|D\|; x) \leq \mu(D; x)$.

1.2. Asymptotic base loci. Following [ELMNP1], one defines the restricted base locus of a big \mathbb{R} -divisor D as

$$\mathbf{B}_-(D) \stackrel{\text{def}}{=} \bigcup_A \mathbf{B}(D + A) ,$$

where the union is taken over all ample divisors A , such that $D + A$ is a \mathbb{Q} -divisor. This locus is a countable union of subvarieties of X by [ELMNP1, Proposition 1.19]

$$\mathbf{B}_-(D) = \bigcup_{m \in \mathbb{N}} \mathbf{B}(D + \frac{1}{m}A) .$$

The augmented base locus of D is defined by

$$\mathbf{B}_+(D) \stackrel{\text{def}}{=} \bigcap_A \mathbf{B}(D - A),$$

where the intersection is taken again over all ample divisors A , such that $D + A$ is a \mathbb{Q} -divisor. It follows quickly from [ELMNP1, Proposition 1.5] that $\mathbf{B}_+(D) = \mathbf{B}(D - \frac{1}{m}A)$ for all $m \gg 0$ and any fixed ample class A .

Proposition 1.2. *Let X be a projective variety, $x \in X$ an arbitrary point. Then*

- (1) $B_+(x) \stackrel{\text{def}}{=} \{\alpha \in N^1(X)_{\mathbb{R}} \mid x \in \mathbf{B}_+(\alpha)\} \subseteq N^1(X)_{\mathbb{R}}$ is closed,
 - (2) $B_-(x) \stackrel{\text{def}}{=} \{\alpha \in N^1(X)_{\mathbb{R}} \mid x \in \mathbf{B}_-(\alpha)\} \subseteq N^1(X)_{\mathbb{R}}$ is open,
- both with respect to the metric topology of $N^1(X)_{\mathbb{R}}$.

For further references and relevant properties of restricted/augmented base loci, we refer the reader to [ELMNP1, KL15], including the proof to Proposition 1.2.

1.3. Newton–Okounkov bodies. Newton–Okounkov bodies have been introduced to projective geometry by Lazarsfeld–Mustață [LM] and Kaveh–Khovanskii [KKh] motivated by earlier work of Okounkov in representation theory [O]. For a big \mathbb{R} -divisor D on X , $\Delta_{Y_\bullet}(D)$ stands for the Newton–Okounkov body of D with respect to the admissible flag Y_\bullet , where

$$Y_\bullet : X = Y_0 \supseteq Y_1 \supseteq \dots \supseteq Y_n$$

is a full flag of (irreducible) subvarieties $Y_i \subseteq X$ with $\text{codim}_X Y_i = i$ and the property that Y_i is smooth at the point Y_n for all $0 \leq i \leq n$. In particular, if X is only assumed to be projective, the center $Y_n = \{x\}$ of an admissible flag must be a smooth point.

Remark 1.3. (Geometry of $\Delta_{Y_\bullet}(D)$) In low dimensions the geometry of $\Delta_{Y_\bullet}(D)$ is well-understood: for curves $\Delta_{Y_\bullet}(D) = [0, \deg D] \subseteq \mathbb{R}$ is a line segment ([LM, Example 1.13]); in the case of surfaces variation of Zariski decomposition [BKS] leads to the fact that Newton–Okounkov bodies are polygons with rational slopes (see [LM, Theorem 6.4] and [KLM, Section 2]).

Note that in dimensions three and above, the situation is no longer purely combinatorial: $\Delta_{Y_\bullet}(D)$ can be non-polyhedral even if D is ample and X is a Mori dream space. At the same time finite generation of the section ring of D ensures the existence of flags with respect to which $\Delta_{Y_\bullet}(D)$ is a rational simplex (see [AKL]).

Next, we quickly recall a few notions and useful facts from [KL15] without proof.

Proposition 1.4 (Equivalent definition of Newton–Okounkov bodies). *Let $\xi \in N^1(X)_{\mathbb{R}}$ be a big \mathbb{R} -class and Y_\bullet be an admissible flag on X . Then*

$$\Delta_{Y_\bullet}(\xi) = \text{closed convex hull of } \{v_{Y_\bullet}(D) \mid D \in \text{Div}_{\geq 0}(X)_{\mathbb{R}}, D \equiv \xi\},$$

where the valuation $v_{Y_\bullet}(D)$, for an effective \mathbb{R} -divisor D , is constructed inductively as in the case of integral divisors.

Proposition 1.5. *Suppose ξ is a big \mathbb{R} -class and Y_\bullet is an admissible flag on X . Then for any $t \in [0, \mu_{Y_1}(\xi))$, we have*

$$\Delta_{Y_\bullet}(\xi)_{v_1 \geq t} = \Delta_{Y_\bullet}(\xi - tY_1) + t\mathbf{e}_1,$$

where $\mathbf{e}_1 = (1, 0, \dots, 0) \in \mathbb{R}^n$.

Lemma 1.6. *Let D be a big \mathbb{R} -divisor and Y_\bullet an admissible flag on X . Then the following hold.*

- (1) *For any real number $\varepsilon > 0$ and any ample \mathbb{R} -divisor A on X , we have $\Delta_{Y_\bullet}(D) \subseteq \Delta_{Y_\bullet}(D + \varepsilon A)$.*
- (2) *If α is an arbitrary nef \mathbb{R} -divisor class, then $\Delta_{Y_\bullet}(D) \subseteq \Delta_{Y_\bullet}(D + \alpha)$.*
- (3) *If α_m is any sequence of nef \mathbb{R} -divisor classes with the property that $\alpha_m - \alpha_{m+1}$ is nef and $\|\alpha_m\| \rightarrow 0$ as $m \rightarrow \infty$ with respect to some norm on $N^1(X)_{\mathbb{R}}$, then*

$$\Delta_{Y_\bullet}(D) = \bigcap_m \Delta_{Y_\bullet}(D + \alpha_m).$$

Definition 1.7. (Valuative points) Let X be a projective variety, Y_\bullet an admissible flag, and D a big \mathbb{Q} -Cartier divisor on X . We call a point $v \in \Delta_{Y_\bullet}(D)$ *valuative*, if it lies in the image of normalized map $\frac{1}{m}v_{Y_\bullet}: |mD| \rightarrow \mathbb{Q}_{\geq 0}$ for some $m \geq 1$, whenever mD becomes Cartier.

Lemma 1.8. *With notation as above, $\text{int } \Delta_{Y_\bullet}(D) \cap \mathbb{Q}^n$ consists of valuative points. If $\Delta_{Y_\bullet}(D)$ contains a small simplex with valuative vertices, then all rational points of the simplex are valuative.*

Proof. Follows from Proposition 1.4 and multiplicative property of v_{Y_\bullet} . □

1.4. Moving Seshadri constants. We recall the necessary information about moving Seshadri constants; our main source is [ELMNP2, Section 6].

Definition 1.9. (Moving Seshadri constant) Let X be a projective variety, $x \in X$ be a smooth point, and D a big \mathbb{R} -divisor with $x \notin B_+(D)$. The *moving Seshadri constant of D at x* is defined as

$$\varepsilon(\|D\|; x) \stackrel{\text{def}}{=} \sup_{f^*D = A + E} \varepsilon(A; x),$$

where the supremum is taken over all projective morphisms $f: Y \rightarrow X$ with Y smooth and f an isomorphism around x , and over all decompositions $f^*D = A + E$, where A is ample, and E is effective with $f^{-1}(x) \notin \text{Supp}(E)$.

If D is nef, then $\varepsilon(\|D\|; x)$ specializes to the usual Seshadri constant $\varepsilon(D; x)$. The formal rules that the moving Seshadri constant obeys can be concisely expressed as follows.

Proposition 1.10. [ELMNP2, Proposition 6.3] *With notation as above, $\varepsilon(\| \cdot \|; x)$ descends to a degree one homogeneous concave function on $\text{Big}(X) \setminus B_+(x)$.*

By virtue of its concavity and the fact that its domain $\text{Big}(X) \setminus B_+(x) \subseteq N^1(X)_{\mathbb{R}}$ is open, $\varepsilon(\| \cdot \|; x)$ is of course a continuous function on it. The highly non-trivial result of [ELMNP2] is that continuity is preserved under extending $\varepsilon(\| \cdot \|; x)$ by zero outside $\text{Big}(X) \setminus B_+(x)$ in $N^1(X)_{\mathbb{R}}$.

Theorem 1.11. [ELMNP2, Theorem 6.2] *Let X be a smooth projective variety, $x \in X$. Then the function $\varepsilon(\|\cdot\|; x) : \mathbf{N}^1(X)_{\mathbb{R}} \rightarrow \mathbb{R}_{\geq 0}$ given by*

$$D \mapsto \begin{cases} \varepsilon(\|D\|; x) & , \text{ if } D \notin B_+(x) \\ 0 & , \text{ otherwise} \end{cases}$$

is continuous.

In Section 5, we offer an alternative extension of $\varepsilon(\|D\|; x)$ over $B_+(x)$.

2. INFINITESIMAL NEWTON–OKOUNKOV BODIES

In this section we define infinitesimal Newton–Okounkov bodies and discuss some of their properties needed in the rest of the paper. Recall that we denote by $\pi : X' \rightarrow X$ the blow-up of X at x with exceptional divisor E . As x is smooth, X' is again a projective variety, and E is an irreducible Cartier divisor on X' , which is smooth as a subvariety of X' .

Definition 2.1. We say that Y_{\bullet} is an *infinitesimal flag over the point x* , if $Y_1 = E$ and each Y_i is a linear subspace in $E \simeq \mathbb{P}^{n-1}$ of dimension $n - i$. We will often write $Y_n = \{z\}$. An *infinitesimal flag over X* is an infinitesimal flag over $x \in X$ for some smooth point x .

The symbol $\tilde{\Delta}_{Y_{\bullet}}(D)$ stands for an infinitesimal Newton–Okounkov body of D , that is,

$$\tilde{\Delta}_{Y_{\bullet}}(D) \stackrel{\text{def}}{=} \Delta_{Y_{\bullet}}(\pi^*D) \subseteq \mathbb{R}_+^n ,$$

where Y_{\bullet} is an infinitesimal flag over x .

Remark 2.2. (Difference in terminology) Note the deviation in terminology from [LM, Section 5.2]; what Lazarsfeld and Mustaa call an infinitesimal Newton–Okounkov body, is in our language (following [KL14]) the generic infinitesimal Newton–Okounkov body.

Remark 2.3. Recently, interesting steps in the infinitesimal direction have been taken by Roe [R].

We start with an observation explaining the shapes of the 'right' kind of simplices that play the role of standard simplices in the infinitesimal theory.

Lemma 2.4. (cf. [KL15, Lemma 3.4]) *Let X be a projective variety, $x \in X$ a smooth point, and A an ample Cartier divisor on X . Then there exists a natural number m_0 such that for any infinitesimal flag Y_{\bullet} over x and for every $m \geq m_0$ there exist global sections $s'_0, \dots, s'_n \in H^0(X', \mathcal{O}_{X'}(\pi^*(mA)))$ for which*

$$v_{Y_{\bullet}}(s'_0) = \mathbf{0} , v_{Y_{\bullet}}(s'_1) = \mathbf{e}_1 , \text{ and } v_{Y_{\bullet}}(s'_i) = \mathbf{e}_1 + \mathbf{e}_i , \text{ for every } 2 \leq i \leq n ,$$

where $\{\mathbf{e}_1, \dots, \mathbf{e}_n\} \subseteq \mathbb{R}^n$ denotes the standard basis.

Proof. The line bundle A is ample, therefore there exists a natural number $m_0 > 0$ such that m_0A is very ample, in particular the linear series $|(m_0 + m)A|$ define embeddings for all $m \geq 0$. As $|m_0A|$ separates tangent directions as well, Bertini's theorem yields the existence of hyperplane

sections $H_1, \dots, H_{n-1} \in |m_0A|$ intersecting transversally at x , and $\tilde{H}_1 \cap \dots \cap \tilde{H}_i \cap E = Y_{i+1}$ for all $i = 1, \dots, n-1$, where \tilde{H}_i denotes the strict transform of H_i through the blow-up map π .

At the same time observe that for any $m \geq m_0$ there exists a global section $t \in H^0(X, \mathcal{O}_X(mA))$ not passing through x . By setting $s'_i \stackrel{\text{def}}{=} \pi^*(t + s_i)$ where $s_i \in H^0(X, \mathcal{O}_X(m_0A))$ is a section associated to H_i , then the sections s'_0, \dots, s'_n satisfy the requirements. \square

Definition 2.5. For a positive real number $\xi \geq 0$, the *inverted standard simplex of size ξ* , denoted by Δ_ξ^{-1} , is the convex hull of the set

$$\Delta_\xi^{-1} \stackrel{\text{def}}{=} \{\mathbf{0}, \xi \mathbf{e}_1, \xi(\mathbf{e}_1 + \mathbf{e}_2), \dots, \xi(\mathbf{e}_1 + \mathbf{e}_n)\} \subseteq \mathbb{R}^n.$$

When $\xi = 0$, then $\Delta_\xi^{-1} = \mathbf{0}$.

A major difference from the non-infinitesimal case is the fact that infinitesimal Newton–Okounkov bodies are also contained in inverted simplices in a very natural way.

Proposition 2.6. *Let D be a big \mathbb{R} -divisor X , then $\Delta_{Y_\bullet}(\pi^*(D)) \subseteq \Delta_{\mu(D;x)}^{-1}$ for any infinitesimal flag Y_\bullet over the point x .*

Proof. By the continuity of Newton–Okounkov bodies inside the big cone it suffices to treat the case when D is a big \mathbb{Q} -divisor. Homogeneity then lets us assume that D is integral. Set $\mu = \mu(D;x)$.

We will follow the line of thought of the proof of [KL14, Proposition 3.2]. Recall that $E \simeq \mathbb{P}^{n-1}$; we will write $[y_1 : \dots : y_n] \in \mathbb{P}^{n-1}$ for a set of homogeneous coordinates in E such that

$$Y_i = \text{Zeroes}(y_1, \dots, y_{i-1}) \subseteq \mathbb{P}^{n-1} = E \text{ for all } 2 \leq i \leq n.$$

With respect to a system of local coordinates (u_1, \dots, u_n) at the point x , the blow-up X' can be described (locally around x) as

$$X' = \{((u_1, \dots, u_n); [y_1 : \dots : y_n]) \mid u_i y_j = u_j y_i \text{ for any } 1 \leq i < j \leq n\}.$$

We can then write a global section s of D in the form

$$s = P_m(u_1, \dots, u_n) + P_{m+1}(u_1, \dots, u_n) + \dots + P_{m+k}(u_1, \dots, u_n)$$

around x , where P_i are homogeneous polynomials of degree i .

We will perform the computation in the open subset $U_n = \{y_n \neq 0\}$, where we can take $y_n = 1$ and the defining equations of the blow-up are given by $u_i = u_n y_i$ for $1 \leq i \leq n-1$. Then

$$s|_{U_n} = u_n^m \cdot (P_m(y_1, \dots, y_{n-1}, 1) + u_n P_{m+1}(y_1, \dots, y_{n-1}, 1) + \dots + u_n^k P_{m+k}(y_1, \dots, y_{n-1}, 1)),$$

in particular, $v_1(s) = m$. Notice that for the rest of $v_i(s)$'s we have to restrict to the exceptional divisor $u_n = 0$ and thus the only term arising in the computation is $P_m(y_1, \dots, y_{n-1}, 1)$.

As $\deg P_m \leq m$, taking into account the algorithm for constructing the valuation vector of a section one can see that indeed

$$v_2(s) + \dots + v_n(s) \leq v_1(s),$$

and this finishes the proof of the proposition. \square

3. RESTRICTED BASE LOCUS VIA NEWTON–OKOUNKOV BODIES

The section is devoted to our characterization of restricted base loci in terms of infinitesimal data. The proofs are variations of those found in [KL15, Section 2].

Theorem 3.1. *Let X be a smooth projective variety, D a big \mathbb{R} -divisor and $x \in X$ an arbitrary point on X . Then the following are equivalent.*

- (1) $x \notin \mathbf{B}_-(D)$.
- (2) There exists an infinitesimal flag Y_\bullet over x such that $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D)$.
- (3) For every infinitesimal flag Y_\bullet over x , one has $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D)$.

Proof. (1) \Rightarrow (3) Assume $x \notin \mathbf{B}_-(D)$, and fix a sequence of ample \mathbb{R} -divisor $(\alpha_m)_{m \in \mathbb{N}}$ so that $\alpha_m - \alpha_{m+1}$ is ample and $D + \alpha_m$ is a \mathbb{Q} -divisor for any $m \geq 1$, and $\|\alpha_m\| \rightarrow 0$ as $m \rightarrow \infty$.

Now, let Y_\bullet be an arbitrary infinitesimal flag over x . Since $x \notin \mathbf{B}_-(D)$, then $x \notin \mathbf{B}(D + \alpha_m)$ for all $m \geq 1$. On the other hand, we have the sequence of equalities

$$\mathbf{B}(\pi^*(D + \alpha_m)) = \pi^{-1}(\mathbf{B}(D + \alpha_m)).$$

In particular, this implies that

$$E \cap \mathbf{B}(\pi^*(D + \alpha_m)) = \emptyset,$$

for all $m \geq 1$. As Y_\bullet is an infinitesimal flag over x , there must exist a sequence of natural numbers $n_m \geq 1$ and a sequence of global sections $s_m \in H^0(X', \mathcal{O}_{X'}(\pi^*(n_m(D + \alpha_m))))$ such that $s_m(z) \neq 0$. This implies that $v_{Y_\bullet}(s_m) = \mathbf{0}$ for each $m \geq 1$. In particular, $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D + \alpha_m)$ for every $m \geq 1$.

Recall that $\pi^* \alpha_m$ is big and semi-ample, therefore

$$\tilde{\Delta}_{Y_\bullet}(D) = \bigcap_{m=1}^{\infty} \tilde{\Delta}_{Y_\bullet}(D + \alpha_m),$$

according to Lemma 1.6, hence $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D)$ as wanted.

The implication (3) \Rightarrow (2) is trivial, and so we are left with checking (2) \Rightarrow (1). Let Y_\bullet be an infinitesimal flag over x so that $\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D)$. Fix an ample \mathbb{R} -divisor A on X and an decreasing sequence of positive real number $(t_m)_{m \in \mathbb{N}}$ such that $\|t_m\| \rightarrow 0$ as $m \rightarrow \infty$, and $D + t_m A$ is a \mathbb{Q} -divisor for all $m \geq 1$. Now, by Lemma 1.6, we know

$$\mathbf{0} \in \tilde{\Delta}_{Y_\bullet}(D) \subseteq \tilde{\Delta}_{Y_\bullet}(D + t_m A)$$

for all $m \geq 0$, therefore $\min \sigma_{\pi^*(D + t_m A)} = 0$ for the sum function $\sigma_{\pi^*(D + t_m A)} : \tilde{\Delta}_{Y_\bullet}(D + t_m A) \rightarrow \mathbb{R}_+$. In particular, this implies, by making use of [KL15, Proposition 2.6], that $\text{mult}_z(\|\pi^*(D + t_m A)\|) = 0$ for all $m \geq 1$, where $Y_n = z$ is the base point of the flag Y_\bullet . Taking into account the string of (in)equalities

$$\text{mult}_x(\|D + t_m A\|) = \text{mult}_E(\|\pi^*(D + t_m A)\|) \leq \text{mult}_z(\|\pi^*(D + t_m A)\|) = 0$$

yields $\text{mult}_x \|D + t_m A\| = 0$ for all $m \geq 1$. As all the divisors $D + t_m A$ were taken to be \mathbb{Q} -divisors, [ELMNP1, Proposition 2.8] leads to $x \notin \mathbf{B}_-(D + t_m A)$ for all $m \geq 1$. But, since

$$\mathbf{B}_-(D) = \bigcup_m \mathbf{B}_-(D + \alpha_m) = \bigcup_m \mathbf{B}(D + \alpha_m)$$

by [ELMNP1, Proposition 1.19], we are done. \square

Remark 3.2. We point out that the implication (1) \Rightarrow (3) remains true under the weaker assumptions that X is a projective variety and $x \in X$ a smooth point. For the converse the answer is unclear since the proof of (2) \Rightarrow (1) uses [ELMNP1, Proposition 2.8], which in turn is verified with the help of multiplier ideals and Nadel vanishing.

Corollary 3.3. *Let X be a smooth projective variety, D a big \mathbb{R} -divisor on X . Then the following are equivalent.*

- (1) D is nef.
- (2) For every point $x \in X$ there exists an infinitesimal flag Y_\bullet over x such that $\mathbf{0} \in \widetilde{\Delta}_{Y_\bullet}(D)$.
- (3) The origin $\mathbf{0} \in \widetilde{\Delta}_{Y_\bullet}(D)$ for every infinitesimal flag over X .

4. AUGMENTED BASE LOCI, INFINITESIMAL NEWTON–OKOUNKOV BODIES, AND JET SEPARATION

In this section, which is the core of the paper, we extend the characterization of augmented base loci in terms of infinitesimal Newton–Okounkov bodies suggested by [KL14, Theorem 3.8] to all dimensions (cf. [KL15, Theorem B] as well). Our statement can be seen as a generalization of Seshadri’s criterion for ampleness. The argument will pass through a study of the connection between infinitesimal Newton–Okounkov bodies and jet separation.

4.1. The main theorem and the largest inverted simplex constant.

Theorem 4.1. *Let X be a smooth projective variety, $x \in X$ an arbitrary (closed) point, D a big \mathbb{R} -divisor on X . Then the following are equivalent.*

- (1) $x \notin \mathbf{B}_+(D)$.
- (2) For every infinitesimal flag Y_\bullet over x there is $\xi > 0$ such that $\Delta_\xi^{-1} \subseteq \widetilde{\Delta}_{Y_\bullet}(D)$.
- (3) There exists an infinitesimal Y_\bullet over x and $\xi > 0$ such that $\Delta_\xi^{-1} \subseteq \widetilde{\Delta}_{Y_\bullet}(D)$.

As an immediate consequence via the equivalence of ampleness and \mathbf{B}_+ being empty, we obtain

Corollary 4.2. *Let X be a smooth projective variety and D a big \mathbb{R} -divisor on X . Then the following are equivalent.*

- (1) D is ample.
- (2) For every point $x \in X$ and every infinitesimal flag Y_\bullet over x there exists a real number $\xi > 0$ for which $\Delta_\xi^{-1} \subseteq \widetilde{\Delta}_{Y_\bullet}(D)$.
- (3) For every point $x \in X$ there exists an infinitesimal flag Y_\bullet over x and a real number $\xi > 0$ such that $\Delta_\xi^{-1} \subseteq \widetilde{\Delta}_{Y_\bullet}(D)$.

We will first give a proof of implication (1) \Rightarrow (2) from Theorem 4.1.

Proposition 4.3. *Let X be a projective variety, $x \in X$ a smooth point, and D a big \mathbb{R} -Cartier divisor on X . If $x \notin \mathbf{B}_+(D)$, then there exists a real number $\xi > 0$ such that $\Delta_\xi^{-1} \subseteq \Delta_{Y_\bullet}(D)$ for any infinitesimal flag Y_\bullet over x*

Proof. This is a modification of the proof of [KL15, Theorem B] using Lemma 2.4; the basic strategy is the same.

Let us first suppose that D is \mathbb{Q} -Cartier. By assumption $x \notin \mathbf{B}_+(D) = \mathbf{B}(D - A)$ for some small ample \mathbb{Q} -Cartier divisor A . Note also that by $\mathbf{B}(\pi^*(D - A)) = \pi^{-1}(\mathbf{B}(D - A))$ this gives

$$\mathbf{B}(\pi^*(D - A)) \cap E = \emptyset$$

as well. Choose a positive integer m large and divisible enough such that $\pi^*(mA)$ becomes integral, satisfies the conclusions of Lemma 2.4, and $\mathbf{B}(\pi^*(D - A)) = \text{Bs}(\pi^*(m(D - A)))$ set-theoretically.

Since $x \notin \text{Bs}(\pi^*(m(D - A)))$, there exists a section $s \in H^0(X', \mathcal{O}_{X'}(\pi^*(mD - mA)))$ with $s(x) \neq 0$, i.e. $v_{Y_\bullet}(s) = \mathbf{0}$. Furthermore, Lemma 2.4 provides global sections $s_0, \dots, s_n \in H^0(X', \mathcal{O}_{X'}(\pi^*(mA)))$ such that $v_{Y_\bullet}(s_0) = \mathbf{0}$, $v_{Y_\bullet}(s_1) = \mathbf{e}_1$ and $v_{Y_\bullet}(s_i) = \mathbf{e}_1 + \mathbf{e}_i$ for all $2 \leq i \leq n$.

Multiplicativity of the valuation map v_{Y_\bullet} then gives

$$v_{Y_\bullet}(s \otimes s_0) = \mathbf{0}, v_{Y_\bullet}(s \otimes s_1) = \mathbf{e}_1 \text{ and } v_{Y_\bullet}(s \otimes s_i) = \mathbf{e}_1 + \mathbf{e}_i \text{ for all } 2 \leq i \leq n.$$

By the construction of Newton–Okounkov bodies, then $\Delta_{1/m}^{-1} \subseteq \tilde{\Delta}_{Y_\bullet}(D)$.

Next, let D be a big \mathbb{R} -divisor for which $x \notin \mathbf{B}_+(D)$, and let A be an ample \mathbb{R} -divisor such that $D - A$ is a \mathbb{Q} -divisor, and $\mathbf{B}_+(D) = \mathbf{B}_+(D - A)$. Then we have $x \notin \mathbf{B}_+(D - A)$, therefore

$$\Delta_\xi^{-1} \subseteq \tilde{\Delta}_{Y_\bullet}(D - A) \subseteq \tilde{\Delta}_{Y_\bullet}(D)$$

for some positive number ξ , according to the \mathbb{Q} -Cartier case and Lemma 1.6. \square

Just as in the surface case, $x \notin \mathbf{B}_+(D)$ implies that $\tilde{\Delta}_{Y_\bullet}(D)$ will contain an inverted standard simplex of some size, hence it makes sense to ask how large these simplices can become (cf. [KL14, Definition 4.5]).

Definition 4.4. (Largest inverted simplex constant) Let X be a projective variety, $x \in X$ a smooth point on X , and D a big \mathbb{R} -divisor with $x \notin \mathbf{B}_-(D)$. For an infinitesimal flag Y_\bullet over x write

$$\xi_{Y_\bullet}(D; x) \stackrel{\text{def}}{=} \sup \left\{ \xi \geq 0 \mid \Delta_\xi^{-1} \subseteq \tilde{\Delta}_{Y_\bullet}(D) \right\}.$$

The *largest inverted simplex constant* $\xi(D; x)$ of D at x is then defined as

$$\xi(D; x) \stackrel{\text{def}}{=} \sup_{Y_\bullet} \xi_{Y_\bullet}(D; x),$$

where Y_\bullet runs through all infinitesimal flags over x . Moreover, if $x \in \mathbf{B}_-(D)$, then let $\xi(D; x) = 0$.

Remark 4.5. As Newton–Okounkov bodies are homogeneous, so is $\xi(\cdot; x)$ as a function on $N^1(X)_{\mathbb{R}}$. Although it is not a priori clear if $\xi(\cdot; x)$ should be continuous, a bit of thought will convince that this is indeed the case over the domain where $x \notin \mathbf{B}_+(D)$.

First, Corollary 4.8 below shows that $\xi_{Y_{\bullet}}(D; x)$ is in fact independent of Y_{\bullet} , therefore we can use one flag for all \mathbb{R} -divisor classes. The natural inclusion

$$\Delta_{Y_{\bullet}}(D) + \Delta_{Y_{\bullet}}(D') \subseteq \Delta_{Y_{\bullet}}(D + D')$$

shows that $\xi(\cdot; x)$ is in fact a concave function on $\text{Big}(X) \setminus \mathbf{B}_+(x)$. This latter is an open subset of $N^1(X)_{\mathbb{R}}$, therefore $\xi(\cdot; x)$ is continuous on its domain. For further results regarding continuity, we advise the reader to look at Corollary 4.12 and Section 5.

Proposition 4.6. *Let X be a normal projective variety, $x \in X$ a smooth point and D a big \mathbb{R} -Cartier divisor on X . Assume that $\Delta_{\xi}^{-1} \subseteq \tilde{\Delta}_{Y_{\bullet}}(D)$ for some infinitesimal flag Y_{\bullet} over x . Then $\Delta_{\xi}^{-1} \subseteq \tilde{\Delta}_{Y'_{\bullet}}(D)$ for all infinitesimal flags Y'_{\bullet} over x .*

Remark 4.7. Normality is used in [FKL, Theorem A], a key ingredient of the proof. The cited result studies the question when the support of an effective \mathbb{R} -divisor is contained in certain augmented base loci in terms of the variation of the volume function.

Proof. The argument below works only for \mathbb{Q} -divisors, passing to the limit delivers the general case (recall that restricted Newton–Okounkov bodies behave in a continuous fashion by [LM, Example 4.22]). Assume that D is a big \mathbb{Q} -divisor on X . For $\xi' \in (0, \xi)$, write $\Delta_{\xi'}^{n-1} \subseteq \mathbb{R}^{n-1}$ for standard simplex of size ξ' and dimension $n - 1$.

Our goal is then to show that

$$\Delta_{Y'_{\bullet}}(\pi^*(D) - \xi'E) \cap \{0\} \times \mathbb{R}^{n-1} = \Delta_{\xi'}^{n-1}$$

for any infinitesimal flag Y'_{\bullet} over x . By continuity it suffices to check this for rational values of ξ' .

So, fix a rational number $\xi' \in (0, \xi)$ and denote by $B \stackrel{\text{def}}{=} \pi^*(D) - \xi'E$. Obviously,

$$\Delta_{Y_{\bullet}}(B + \lambda E) = \Delta_{Y_{\bullet}}(\pi^*D - (\xi' - \lambda)E)$$

for any $\lambda < \xi'$. The condition $\Delta_{\xi}^{-1} \subseteq \tilde{\Delta}_{Y_{\bullet}}(D)$ and Proposition 1.5 imply

$$\text{vol}_{\mathbb{R}^n}(\Delta_{Y_{\bullet}}(B + \lambda E)) > \text{vol}_{\mathbb{R}^n}(\Delta_{Y_{\bullet}}(B))$$

for any rational number $0 < \lambda < \xi'$. Then [LM, Theorem A] gives $\text{vol}_X(B + \lambda E) > \text{vol}_X(B)$, which, via [FKL, Theorem B] leads to $E \notin \mathbf{B}_+(B)$.

The significance of this condition is that it grants us access to the slicing theorem [LM, Theorem 4.24]. In particular,

$$\Delta_{Y_{\bullet}|E}(B) = \Delta_{Y_{\bullet}}(B)|_{x_1=0} = \Delta_{\xi'}^{n-1},$$

where left-hand side denotes the appropriate restricted Newton–Okounkov body (see [LM, (2.7)]). By the same token, since $E \notin \mathbf{B}_+(B)$, we have

$$\text{vol}_{X|E}(B) = (n-1)! \text{vol}_{\mathbb{R}^{n-1}}(\Delta_{Y_{\bullet}|E}(B)) = (n-1)! \text{vol}_{\mathbb{R}^{n-1}}(\Delta_{\xi'}^{n-1}).$$

Note that both extremes are independent of the choice of the flag, hence we have

$$\mathrm{vol}_{\mathbb{R}^{n-1}}(\Delta_{Y'_\bullet|E}(B)) = \mathrm{vol}_{\mathbb{R}^{n-1}}(\Delta_{\xi'}^{n-1})$$

for any infinitesimal flag Y'_\bullet on X' .

It follows from Proposition 2.6 that

$$\Delta_{Y'_\bullet|E}(B) = \Delta_{Y'_\bullet}(B)|_{x_1=0} \subseteq \Delta_{\xi'}^{n-1},$$

however, as the two convex bodies have equal volume, they must coincide. This means that $\Delta_{Y'_\bullet}(B)|_{x_1=0} = \Delta_{\xi'}^{n-1}$ as required. \square

Corollary 4.8. *With notation as above, $\xi(D;x) = \xi_{Y'_\bullet}(D;x)$ for all infinitesimal flags Y'_\bullet over x .*

4.2. Inverted standard simplices and jet separation. Arguably one of the most important ingredients of the proof of Theorem 4.1 is the following connection between infinitesimal Newton–Okounkov bodies and jet separation of adjoint bundles.

Proposition 4.9. *(Infinitesimal Newton–Okounkov bodies and jet separation) Let X be an n -dimensional smooth projective variety, D a big Cartier divisor, and x be a (closed) point on X . Assume that there exists a positive real number ε and a natural number k with the property that $\Delta_{n+k+\varepsilon}^{-1} \subseteq \Delta_{Y'_\bullet}(\pi^*(D))$ for every infinitesimal flag Y'_\bullet over x . Then $K_X + D$ separates k -jets.*

Proof. By definition (see [D], also [PAG1, Definiton 5.1.15] and [PAG1, Proposition 5.1.19]), what we need to prove is that the restriction map

$$H^0(X, \mathcal{O}_X(K_X + D)) \longrightarrow H^0(X, \mathcal{O}_X(K_X + D) \otimes \mathcal{O}_{X,x}/\mathfrak{m}_{X,x}^{k+1})$$

is surjective.

Transferring the question to the blow-up X' , this is equivalent to requiring

$$(4.9.1) \quad H^0(X, \mathcal{O}_{X'}(\pi^*(K_X + D))) \rightarrow H^0(X', \mathcal{O}_{X'}(\pi^*(K_X + D)) \otimes \mathcal{O}_{X'}/\mathcal{O}_{X'}(-(k+1)E))$$

to be surjective.

In order to do check surjectivity in (4.9.1), let us write $B \stackrel{\mathrm{def}}{=} \pi^*(D) - (n+k)E$. By Proposition 1.5, we have

$$\Delta_{Y'_\bullet}(B) = \Delta_{Y'_\bullet}(\pi^*(D))_{x_0 \geq n+k} - (n+k, 0, \dots, 0)$$

for any infinitesimal flag Y'_\bullet over the point x . In particular, B is a big line bundle with the property that the origin $\mathbf{0} \in \Delta_{Y'_\bullet}(B)$ for any infinitesimal flag Y'_\bullet . As a consequence of Theorem 3.1, we obtain that $\mathbf{B}_-(B) \cap E = \emptyset$. Thus $\mathrm{Zeroes}(\mathcal{I}(X', \|B\|)) \cap E = \emptyset$ via [ELMNP1, Corollary 10].

To finish off the proof, we will make use of a variant of the classical argument to deduce the required surjectivity. Recall that $B = \pi^*D - (n+k)E$, and $K_{X'} = \pi^*K_X + (n-1)E$, therefore we have the short exact sequence

$$0 \rightarrow \mathcal{O}_{X'}(K_{X'} + B) \otimes \mathcal{I}(X', \|B\|) \rightarrow \mathcal{O}_{X'}(\pi^*(K_X + D)) \rightarrow \mathcal{O}_{X'}(\pi^*(K_X + D)) \otimes (\mathcal{L} \oplus \mathcal{O}_{(k+1)E}) \rightarrow 0,$$

where \mathcal{L} stands for the structure sheaf determined by the closed subscheme associated to the ideal $\mathcal{I}(X', \|B\|)$; note that this latter has support disjoint from E .

Since B is a big line bundle, by Nadel's vanishing for asymptotic multiplier ideals [PAG2, Theorem 11.2.12.(ii)] we have

$$H^1(X', \mathcal{O}_{X'}(K_{X'} + B) \otimes \mathcal{I}(X', \|B\|)) = 0 ,$$

therefore the restriction map

$$H^0(X', \mathcal{O}_{X'}(\pi^*(K_X + D))) \longrightarrow H^0(X', \mathcal{O}_{X'}(\pi^*(K_X + D)) \otimes (\mathcal{L} \oplus \mathcal{O}_{(k+1)E}))$$

is surjective, but then so is

$$H^0(X', \mathcal{O}_{X'}(\pi^*(K_X + D))) \longrightarrow H^0(X', \mathcal{O}_{X'}(\pi^*(K_X + D)) \otimes \mathcal{O}_{(k+1)E}) ,$$

as required. \square

Now we are in a position to finish the proof of Theorem 4.1; our main tool is going to be the connection between moving Seshadri constants and largest inverted simplex constants via jet separation (cf. [ELMNP2, Proposition 6.6])

Proposition 4.10. *Let D be a big \mathbb{R} -divisor on a smooth projective variety X and $x \in X$ a closed point. If $\xi(D; x) > 0$, then $\xi(D; x) = \varepsilon(\|D\|; x)$.*

Proof. Let us first assume that D is a big \mathbb{Q} -divisor; we wish to show that

$$(4.10.2) \quad \xi(D; x) = \limsup_{m \rightarrow \infty} \frac{s(mD; x)}{m} = \varepsilon(\|D\|; x) ,$$

where the latter equality is [ELMNP2, Proposition 6.6]. Then one can go on and use (4.10.2) and Proposition 4.9 to deduce $\xi(D; x) = \varepsilon(\|D\|; x)$.

Our first goal is to check $\varepsilon(\|D\|; x) \geq \xi(D; x)$. Since both expressions are homogeneous, it will suffice to show $\varepsilon(\|D\|; x) \geq n$ whenever $\xi(D; x) > n$. Let $r > 0$ be a natural number so that rD becomes integral. Then, by homogeneity, $\xi(mrD; x) > mnr$, and Proposition 4.9 gives

$$s(K_X + mrD; x) \geq mnr - n .$$

Consequently, by taking multiples we obtain

$$\frac{s(k(K_X + mrD); x)}{k} \geq mnr - n, \text{ for any } m, k \geq 1 ,$$

in particular, by [ELMNP2, Proposition 6.6] one has

$$\varepsilon(\|K_X + mrD\|; x) = \limsup_{k \rightarrow \infty} \frac{s(k(K_X + mrD); x)}{k} \geq mnr - n .$$

On the other hand, [ELMNP2, Theorem 6.2] says that the function $N^1(X)_{\mathbb{R}} \ni \alpha \mapsto \varepsilon(\|\alpha\|; x) \in \mathbb{R}_+$ is continuous, therefore

$$\varepsilon(\|D\|; x) = \frac{1}{r} \limsup_{m \rightarrow \infty} \frac{\varepsilon(\|K_X + mrD\|; x)}{m} \geq n .$$

For the converse inequality $\varepsilon(\|D\|;x) \leq \xi(D;x)$, we will show that whenever D is an integral divisor separating s -jets at the point x , then $\Delta_s^{-1} \subseteq \Delta_{Y_\bullet}(\pi^*(D))$ for any infinitesimal flag Y_\bullet over x . Note that Proposition 4.6 shows that it suffices check this for one such flag.

To this end, choose a system of local coordinates $\{u_1, \dots, u_n\}$ at x and choose the infinitesimal flag Y_\bullet in such a way that each Y_{i+1} is given by the intersection of E with the proper transforms of u_1, \dots, u_i . Because D separates s -jets at x , there exist sections $s_1, \dots, s_n \in H^0(X, \mathcal{O}_X(D))$ such that $s_i = u_i^s$ locally. Analogously to the proof of Lemma 2.4, we see that $v_{Y_\bullet}(\pi^*(s_1)) = s \cdot \mathbf{e}_1$ and $v_{Y_\bullet}(\pi^*(s_i)) = s \cdot (\mathbf{e}_1 + \mathbf{e}_i)$. The origin is contained in $\tilde{\Delta}_{Y_\bullet}(D)$ since $\xi(D;x) > 0$.

Lastly, it remains to deal with the case when D is a big \mathbb{R} -divisor, which we will do by reduction to the rational case. Fix a sequence of ample \mathbb{R} -divisors $(\alpha_l)_{l \in \mathbb{N}}$ for which $\lim_{l \rightarrow \infty} \|\alpha_l\| = 0$ for an arbitrary norm on $N^1(X)_{\mathbb{R}}$, $D + \alpha_l$ is a \mathbb{Q} -divisor, and $\alpha_{l+1} - \alpha_l$ is ample for any $l \geq 1$.

Then Lemma 1.6 yields

$$\Delta_{Y_\bullet}(\pi^*(D)) = \bigcap_{l \in \mathbb{N}} \Delta_{Y_\bullet}(\pi^*(D + \alpha_l))$$

for any infinitesimal flag Y_\bullet over x . As a consequence, $\lim_{l \rightarrow \infty} \xi(D + \alpha_l; x) = \xi(D; x)$; however, since each class $D + \alpha_l$ is a \mathbb{Q} -divisor, we know that $\xi(D + \alpha_l) = \varepsilon(\|D + \alpha_l\|; x)$ for any $l \in \mathbb{N}$. Continuity of moving Seshadri constants [ELMNP2, Theorem 6.2] then concludes the proof. \square

Proof of Theorem 4.1. The implication (1) \Rightarrow (2) has been taken care of in Proposition 4.3, as (2) \Rightarrow (3) is formal, we are left with (3) \Rightarrow (1). However, if there exists an infinitesimal flag Y_\bullet over x with an inverted standard simplex contained in it, then $\xi(D; x) > 0$, hence Proposition 4.10 yields $\varepsilon(\|D\|; x) = \xi(D; x) > 0$, which by definition means $x \notin \mathbf{B}_+(D)$. \square

We obtain a sequence of interesting corollaries.

Corollary 4.11. *Let D be a big \mathbb{R} -divisor on a smooth projective variety X . Then*

$$\xi(D; x) = \varepsilon(\|D\|; x)$$

for any (closed) point $x \in X$.

Proof. If $x \notin \mathbf{B}_+(D)$, then this is immediate from Theorem 4.1 and Proposition 4.10. If $x \in \mathbf{B}_+(D) \setminus \mathbf{B}_-(D)$, then $\xi(D; x) = 0$ by Proposition 4.6 and $\varepsilon(\|D\|; x) = 0$ by definition. In the last case $x \in \mathbf{B}_-(D)$, both invariants are zero by definition. \square

Corollary 4.12. *For a smooth projective variety X and a point $x \in X$, the function*

$$\begin{aligned} \xi(\cdot; x): N^1(X)_{\mathbb{R}} &\longrightarrow \mathbb{R}_+ \\ D &\longmapsto \xi(D; x) \end{aligned}$$

is continuous.

Proof. Follows easily from Corollary 4.11 and [ELMNP2, Theorem 6.2]. \square

Corollary 4.13. *Let D be a big \mathbb{Q} -divisor on X , $x \in X$ a closed point, and Y_\bullet an infinitesimal flag over x . If $\Delta_\xi^{-1} \subseteq \Delta_{Y_\bullet}(\pi^*(D))$ for some $\xi > 0$, then all vectors in $\Delta_\xi^{-1} \cap \mathbb{Q}^n$ not lying on the face generated by the points $\lambda \cdot \mathbf{e}_1, \lambda(\mathbf{e}_1 + \mathbf{e}_2), \dots, \lambda(\mathbf{e}_1 + \mathbf{e}_n)$ are valuate.*

Proof. This is a consequence of Lemma 1.8, the inequality

$$\lim_{k \rightarrow \infty} \frac{s(kD; x)}{k} \geq \lambda ,$$

and the definition of jet separation constants. One can see by the proof of Proposition 4.10 all vectors with rational coordinates sitting on one of the rays coming out of the origin in the inverted simplex come from a basis for some power of the maximal ideal of x . \square

5. THE EXTENDED SESHADRI FUNCTION

In this section we briefly discuss an extension of moving Seshadri constants completing in some ways the picture considered in [ELMNP2]. We also give an example where the Seshadri constant function inside the ample cone is not everywhere differentiable.

First, recall the notion of asymptotic multiplicity: for a point $x \in X$ on a smooth projective variety X , the *asymptotic multiplicity* of a big \mathbb{R} -divisor D is defined as

$$\text{mult}_x \|D\| \stackrel{\text{def}}{=} \inf_{D'} \{ \text{mult}_x(D') \} ,$$

where the minimum is over all effective \mathbb{R} -divisors with $D' \equiv D$ (see [ELMNP1] for the general theory).

Note that $\text{mult}_x \|D\| > 0$ precisely when $x \in \mathbf{B}_-(D)$ by [ELMNP1, Proposition 2.9]; in contrast with the various largest simplex constants and the geometric definition of the moving Seshadri constant, $\text{mult}_x \|D\|$ concerns the situation when the point $x \in \mathbf{B}_-(D)$. Our goal is to see this invariant through the eyes of infinitesimal Newton–Okounkov bodies, and use this relation to connect asymptotic multiplicities to moving Seshadri constants.

Proposition 5.1. *Let D be an \mathbb{R} -divisor on X , $x \in \mathbf{B}_-(D)$, and denote by $r \stackrel{\text{def}}{=} \text{mult}_x \|D\|$. Then for any infinitesimal flags Y_\bullet over the point x , the following hold*

- (1) $\tilde{\Delta}_{Y_\bullet}(D) \subseteq r \cdot \mathbf{e}_1 + \mathbb{R}_+^n$. In particular, $E \subseteq \mathbf{B}_+(\pi^*(D) - rE)$.
- (2) $E \not\subseteq \mathbf{B}_-(\pi^*(D) - rE)$. In particular $\tilde{\Delta}_{Y_\bullet}(D) \cap \{r\} \times \mathbb{R}^{n-1} \neq \emptyset$.
- (3) The intersection $\tilde{\Delta}_{Y_\bullet}(D) \cap \{r\} \times \mathbb{R}^{n-1}$ has empty interior in \mathbb{R}^{n-1} .

Proof. (1) As $x \in \mathbf{B}_-(D)$, the asymptotic multiplicity $r = \text{mult}_x \|D\|$ is strictly positive. By the definition of asymptotic multiplicity coupled with the fact that $\text{mult}_x(D') = \text{ord}_E(\pi^*(D')) = v_1(\pi^*(D'))$ for any effective \mathbb{R} -divisor $D' \equiv D$ we obtain, $\Delta_{Y_\bullet}(\pi^*(D)) \subseteq r \cdot \mathbf{e}_1 + \mathbb{R}_+^n$.

Take an arbitrary point $z \in E$ and an infinitesimal flag Y_\bullet centered at $z \in X'$. Proposition 1.5 implies that $\mathbf{0} \notin \Delta_{Y_\bullet}(\pi^*D - tE)$ for $0 \leq t < r$. Then $z \in \mathbf{B}_-(\pi^*D - tE) \subseteq \mathbf{B}_+(\pi^*D - tE)$ follows from [KL15, Theorem A] for all $0 \leq t < r$. Using Proposition 1.2.(i), then we know that $B_+(z)$ is closed in the big cone and in particular this yields that $z \in \mathbf{B}_+(\pi^*D - rE)$ as well.

(2) Observe that $\pi^*D - rE$ is big (it has the same volume as D has), therefore $\pi^*D - (r+t)E$ is big for all $0 < t \ll 1$. By the definition of asymptotic multiplicity, $\text{mult}_E \|\pi^*D - (r+t)E\| = 0$ for all $0 < t \ll 1$, in particular $E \not\subseteq \mathbf{B}_-(\pi^*D - (r+t)E)$. But then $z \notin \mathbf{B}_-(\pi^*D - (r+t)E)$ for all

rational values $0 < t \ll 1$ provided $z \in E$ is very general. Now, making use of Proposition 1.2.(ii) we know that $\text{Big}(X')_{\mathbb{R}} \setminus B_-(z)$ is closed. In particular, this yields that $z \notin \mathbf{B}_-(\pi^*D - rE)$.

(3) Let us first point out that $E \not\subseteq \mathbf{B}_+(\pi^*(D) - (r+t)E)$ for any $0 < t \ll 1$. To see this, recall that by (2) above, $\tilde{\Delta}_{Y_\bullet}(D) \cap \{r\} \times \mathbb{R}^{n-1} \neq \emptyset$. Second, $\tilde{\Delta}_{Y_\bullet}(D)$ is a full dimensional convex body, therefore

$$\text{vol}_{X'}(\pi^*(D) - rE) > \text{vol}_{X'}(\pi^*(D) - (t+r)E),$$

by [LM, Theorem A]. But then [FKL, Theorem A] gives $E \not\subseteq \mathbf{B}_+(\pi^*(D) - (r+t)E)$ for any $0 < t \ll 1$.

To finish the proof, suppose for a contradiction that

$$\text{vol}_{\mathbb{R}^{n-1}}(\tilde{\Delta}_{Y_\bullet}(D) \cap \{r\} \times \mathbb{R}^{n-1}) > 0.$$

By the slicing theorem [LM, Theorem 4.24] and the fact that $E \not\subseteq \mathbf{B}_+(\pi^*(D) - (r+t)E)$ for any $0 < t \ll 1$, we obtain

$$\lim_{t \rightarrow 0} \left(\text{vol}_{X'|E}(\pi^*(D) - (t+r)E) \right) \geq \text{vol}_{\mathbb{R}^{n-1}}(\tilde{\Delta}_{Y_\bullet}(D) \cap \{r\} \times \mathbb{R}^{n-1}) > 0.$$

On the other hand, [ELMNP2, Theorem 5.7] forces the limit on the left-hand side to be zero, since E is an irreducible component of $\mathbf{B}_+(\pi^*(D) - rE)$ by (1), a contradiction. \square

Lemma 5.2. *Let $(D_k)_{k \in \mathbb{N}}$ be a sequence of big \mathbb{R} -divisors on a smooth projective variety X converging to a big \mathbb{R} -divisor D , let $x \in X$ be a point. Then*

- (1) *If $\varepsilon(\|D_k\|; x) > 0$ for all $k \in \mathbb{N}$, and $\lim_{k \rightarrow \infty} \varepsilon(\|D_k\|; x) = 0$, then $x \in \mathbf{B}_+(D) \setminus \mathbf{B}_-(D)$.*
- (2) *If $\text{mult}_x(\|D_k\|) > 0$ for all $k \in \mathbb{N}$, and $\lim_{k \rightarrow \infty} \text{mult}_x(\|D_k\|) = 0$, then $x \in \mathbf{B}_+(D) \setminus \mathbf{B}_-(D)$.*

Proof. (1) By Corollary 4.11 it is legal to write $\xi_k \stackrel{\text{def}}{=} \xi(\|D_k\|; x) = \varepsilon(\|D_k\|; x)$ for each $k \in \mathbb{N}$. Fixing an infinitesimal flag Y_\bullet over x , by definition we have $\mathbf{0} \in \Delta_{\xi_k}^{-1} \subseteq \tilde{\Delta}_{Y_\bullet}(D_k)$. By continuity of Newton–Okounkov bodies we obtain $\mathbf{0} \in \Delta_{Y_\bullet}(\pi^*(D))$, we can conclude by Theorem 3.1, $x \notin \mathbf{B}_-(D)$.

On the other hand, $x \in \mathbf{B}_+(D)$ follows from the continuity of the moving Seshadri constant as a function on the Néron-Severi space.

(2) Since $\text{mult}_x(\|D_k\|) > 0$, [ELMNP1, Theorem B] implies that $x \in \mathbf{B}_-(D_k)$ for all $k \in \mathbb{N}$. But then $x \in \mathbf{B}_+(D_k)$ for all $k \in \mathbb{N}$ as well, whence $x \in \mathbf{B}_+(D)$ according to [KL15, Proposition 1.2]. For $x \notin \mathbf{B}_-(D)$ note that asymptotic multiplicity is continuous on the big cone (see [ELMNP1, Theorem A]), therefore $\text{mult}_x(\|D\|) = 0$, and consequently $x \notin \mathbf{B}_-(D)$. \square

By Corollary 4.11, and Lemma 5.2 one can glue the functions $\varepsilon(\|\cdot\|; x)$ and $-\text{mult}_x(\|\cdot\|)$ giving rise to a continuous extension of the moving Seshadri constant function which is nowhere zero on the open subset $B_-(x) \subseteq \mathbf{N}^1(X)_{\mathbb{R}}$.

Definition 5.3 (Extended Seshadri function). Let X be a smooth projective variety, $x \in X$. We define the (extended) Seshadri function $\varepsilon_x: \text{Big}(X) \rightarrow \mathbb{R}_{\geq 0}$ associated to the point $x \in X$ by

$$\varepsilon_x(D) \stackrel{\text{def}}{=} \begin{cases} \varepsilon(\|D\|; x) & \text{if } D \notin B_+(x) \\ 0 & \text{if } D \in B_+(x) \setminus B_-(x) \\ -\text{mult}_x \|D\| & \text{if } D \in B_-(x). \end{cases}$$

Remark 5.4. Since both the asymptotic multiplicity and the moving Seshadri constant are concave on the domain where they are non-zero, it is not unnatural to hope that the extended Seshadri function will retain this property. We shall see that this is indeed the case in the example below.

We end this section with an explicit computation of the extended Seshadri function; an interesting feature of the example is that ε_x is not everywhere differentiable even inside the ample cone.

Example 5.5 (A non-differentiable Seshadri function). Let $p \in \mathbb{P}^2$ be a point and denote by $\pi_1: X \stackrel{\text{def}}{=} \text{Bl}_p(\mathbb{P}^2) \rightarrow \mathbb{P}^2$ the blow-up of \mathbb{P}^2 at the point p with exceptional divisor E . We pick a point $x \in E$, and then pursue to compute the function ε_x on the pseudo-effective cone $\overline{\text{Eff}}(X) = \mathbb{R}_+ E + \mathbb{R}_+(H - E)$, where H is the pullback of the class of a line.

The function ε_x being homogeneous of degree one, it will suffice to determine the values of ε_x as we traverse the line segment $[E, H - E] \subseteq N^1(X)_{\mathbb{R}} = \mathbb{R}^2$. To this end, set

$$F_t \stackrel{\text{def}}{=} tH + (1 - 2t)E \quad \text{for all } 0 \leq t \leq 1.$$

Observe that for $t \in [0, \frac{1}{2})$ we have $x \in \mathbf{B}_-(F_t)$, and

$$\varepsilon_x(F_t) = -\text{mult}_x \|tH + (1 - 2t)E\| = 2t - 1.$$

If $\frac{1}{2} \leq t \leq 1$, then F_t is nef, hence $\varepsilon_x(F_t) = \varepsilon(\|F_t\|; x) = \varepsilon(F_t; x)$. The Seshadri constants $\varepsilon(F_t; x)$ are somewhat more complicated to compute, this will take up the remaining part of our example.

Thus, let $\pi_2: X' \rightarrow X$ denote the blow-up of X at the point x . Write $\pi = \pi_2 \circ \pi_1$ for the composition of the two blow-ups. On X' we have precisely three negative curves:

- E_1 = the strict transform of the exceptional divisor of π_1 under π_2 ,
- E_2 = the exceptional divisor of the blow-up π_2 ,
- E_3 = the strict transform of the line of class $H - E$ on X going through the point x .

The intersection matrix of the curves E_i is

$$(E_i \cdot E_j)_{1 \leq i \leq 3, 1 \leq j \leq 3} = \begin{pmatrix} -2 & 1 & 0 \\ 1 & -1 & 1 \\ 0 & 1 & -1 \end{pmatrix}.$$

In the basis (E_1, E_2, E_3) of $N^1(X')_{\mathbb{R}}$, the hyperplane class H' is given as

$$H' = E_1 + 2E_2 + E_3.$$

Along with H' , the divisors $H' + E_3$ and $E_2 + E_3$ turn out to be nef as well, and the three generate the nef cone of X' . In this notation,

$$D_t \stackrel{\text{def}}{=} \pi_2^* F_t = \pi_2^*(tH + (1-2t)E) = tH' + (1-2t)(E_1 + E_2),$$

which can in turn be written in the form

$$D_t = (1-t)H' + (2t-1)(E_2 + E_3) \text{ for all } 1/2 \leq t \leq 1.$$

This means in particular that D_t sits on the face of the nef cone generated by H' and $E_2 + E_3$ for all $1/2 \leq t \leq 1$.

As one can check, that the ray $D_t - \varepsilon E_1$ leave the nef cone through the the face generated by the divisors H' and $H' + E_3$ whenever $t \in [\frac{1}{2}, \frac{2}{3}]$, and throught the face generated by the divisors $H' + E_3$ and $E_1 + E_3$ for $t \in [\frac{2}{3}, 1]$.

As a result, ε_x is going to be piecewise linear, and it is not going to be differentiable at $t = \frac{2}{3}$. The full computation goes as follows.

For $t \in [\frac{1}{2}, \frac{2}{3}]$, the ray $D_t - \varepsilon E_1$ hits the boundary of the nef cone at $\varepsilon = 2t - 1$, in the divisor

$$D_t - (2t-1)E_1 = (2-3t)H' + (2t-1)(H' + E_3).$$

In particular, $\varepsilon_x(F_t) = 2t - 1$ on the interval $[\frac{1}{2}, \frac{2}{3}]$.

On the other hand, if $t \in [\frac{2}{3}, 1]$, the ray $D_t - \varepsilon E_1$ reaches the boundary of the nef cone at $\varepsilon = 1 - t$, in the divisor

$$D_t - (1-t)E_1 = (1-t)(H' + E_3) + (3t-2)(E_1 + E_3),$$

and we obtain $\varepsilon_x(t) = 1 - t$ on the interval $[\frac{2}{3}, 1]$. Putting all this together, the Seshadri function on the line segment $[E, H - E]$ is given by

$$\varepsilon_x(F_t) = \begin{cases} 2t - 1 & \text{if } t \in [0, \frac{1}{2}] \\ 2t - 1 & \text{if } t \in [\frac{1}{2}, \frac{2}{3}] \\ 1 - t & \text{if } t \in [\frac{2}{3}, 1] \end{cases}.$$

REFERENCES

- [AKL] Dave Anderson, Alex Küronya, and Victor Lozovanu, *Okounkov bodies of finitely generated divisors*, International Mathematics Research Notices **132** (2013), no. 5, 1205–1221. [↑\(document\)](#), [1.3](#)
- [BKS] Thomas Bauer, Alex Küronya, and Tomasz Szemberg, *Zariski decompositions, volumes, and stable base loci*, Journal für die reine und angewandte Mathematik **576** (2004), 209–233. [↑1.3](#)
- [B] Sébastien Boucksom, *Corps D'Okounkov*, Séminaire Bourbaki **65** (Unknown Month 2012), no. 1059, 1–38. [↑\(document\)](#)
- [BCL] S. Boucksom, S. Cacciola, and A. F. Lopez, *Augmented base loci and restricted volumes on normal varieties* (2013). arXiv:1305.4284. [↑\(document\)](#)
- [CHPW] Sung Rak Choi, Yoonsuk Hyun, Jinhyung Park, and Joonyeong Won, *Asymptotic base loci via Okounkov bodies* (2015). arXiv:1506.00817. [↑\(document\)](#)
- [D] Jean-Pierre Demailly, *Singular Hermitian metrics on positive line bundles*, Complex Algebraic Varieties (Bayreuth, 1990), 1992, pp. 87–104. Lect. Notes in Math., vol. 1507. [↑\(document\)](#), [4.2](#)

- [ELMNP1] Lawrence Ein, Robert Lazarsfeld, Mircea Mustață, Michael Nakamaye, and Mihnea Popa, *Asymptotic invariants of base loci*, Ann. Inst. Fourier (Grenoble) **56** (2006), no. 6, 1701–1734. ↑[1.2](#), [1.2](#), [3](#), [3.2](#), [4.2](#), [5](#), [5](#)
- [ELMNP2] ———, *Restricted volumes and base loci of linear series*, Amer. J. Math. **131** (2009), no. 3, 607–651, DOI 10.1353/ajm.0.0054. ↑[\(document\)](#), [1.4](#), [1.10](#), [1.4](#), [1.11](#), [4.2](#), [4.2](#), [4.2](#), [5](#), [5](#)
- [FKL] Mihai Fulger, János Kollár, and Brian Lehmann, *Volume and Hilbert function of \mathbb{R} -divisors* (2015). arXiv:1503.02324. ↑[\(document\)](#), [4.7](#), [4.1](#), [5](#)
- [J] Shin-Yao Jow, *Okounkov bodies and restricted volumes along very general curves*, Adv. Math. **223** (2010), no. 4, 1356–1371, DOI 10.1016/j.aim.2009.09.015. ↑[\(document\)](#)
- [KKh] Kiumars Kaveh and A. G. Khovanskii, *Newton-Okounkov bodies, semigroups of integral points, graded algebras and intersection theory*, Ann. of Math. (2) **176** (2012), no. 2, 925–978, DOI 10.4007/annals.2012.176.2.5. ↑[\(document\)](#), [1.3](#)
- [K] Alex Küronya, *Positivity of restrictions on subvarieties and vanishing of higher cohomology*, Annales de l’Institut Fourier **63** (2013), no. 5, 1717–1737. ↑[\(document\)](#)
- [KL14] Alex Küronya and Victor Lozovanu, *Local positivity of linear series on surfaces*. arXiv:1411.6205. ↑[\(document\)](#), [2.2](#), [2](#), [4](#), [4.1](#)
- [KL15] ———, *Positivity of line bundles and Newton–Okounkov bodies*. arXiv:1506.06525. ↑[\(document\)](#), [1.2](#), [1.3](#), [2.4](#), [3](#), [3](#), [4](#), [4.1](#), [5](#), [5](#)
- [KLM] Alex Küronya, Victor Lozovanu, and Catriona Maclean, *Convex bodies appearing as Okounkov bodies of divisors*, Adv. Math. **229** (2012), no. 5, 2622–2639, DOI 10.1016/j.aim.2012.01.013. ↑[\(document\)](#), [1.3](#)
- [LM] Robert Lazarsfeld and Mircea Mustață, *Convex bodies associated to linear series*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 5, 783–835. ↑[\(document\)](#), [1.3](#), [1.3](#), [2.2](#), [4.1](#), [5](#)
- [LSS] Patrycja Łuszcz-Świdecka and David Schmitz, *Minkowski decomposition of Okounkov bodies on surfaces*, J. Algebra **414** (2014), 159–174. ↑[\(document\)](#)
- [N] Michael Nakamaye, *Base loci of linear series are numerically determined*, Trans. Amer. Math. Soc. **355** (2002), 551–566. ↑[\(document\)](#)
- [PAG1] Robert Lazarsfeld, *Positivity in algebraic geometry. I*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 48, Springer-Verlag, Berlin, 2004. ↑[\(document\)](#), [4.2](#)
- [PAG2] ———, *Positivity in algebraic geometry. II*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge., vol. 49, Springer-Verlag, Berlin, 2004. ↑[4.2](#)
- [O] Andrei Okounkov, *Brunn-Minkowski inequality for multiplicities*, Invent. Math. **125** (1996), no. 3, 405–411. ↑[1.3](#)
- [PSU] Piotr Pokora, David Schmitz, and Stefano Urbinati, *Minkowski bases on toric varieties* (2014). ↑[\(document\)](#)
- [R] Joaquim Roé, *Local positivity in terms of Newton–Okounkov bodies* (2015). arXiv:1505.02051. ↑[2.3](#)

ALEX KÜRONYA, JOHANN-WOLFGANG-GOETHE UNIVERSITÄT FRANKFURT, INSTITUT FÜR MATHEMATIK,
ROBERT-MAYER-STRASSE 6-10., D-60325 FRANKFURT AM MAIN, GERMANY

BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, DEPARTMENT OF ALGEBRA, EGRY JÓZSEF U. 1.,
H-1111 BUDAPEST, HUNGARY

E-mail address: kuronya@math.uni-frankfurt.de

VICTOR LOZOVANU, UNIVERSITÀ DEGLI STUDI DI MILANO–BICOCCA, DIPARTIMENTO DI MATEMATICA E
APPLICAZIONI, VIA COZZI 53,, I-20125 MILANO, ITALY

E-mail address: victor.lozovanu@unimib.it