

Genetics of lipedema: new perspectives on genetic research and molecular diagnoses

S. PAOLACCI¹, V. PRECONE², F. ACQUAVIVA³, P. CHIURAZZI^{4,5}, E. FULCHERI^{6,7}, M. PINELLI^{3,8}, F. BUFFELLI^{9,10}, S. MICHELINI¹¹, K.L. HERBST¹², V. UNFER¹³, M. BERTELLI²; GNEOB PROJECT

¹MAGI'S LAB, Rovereto (TN), Italy

²MAGI EUREGIO, Bolzano, Italy

³Department of Translational Medicine, Section of Pediatrics, Federico II University, Naples, Italy

⁴Istituto di Medicina Genomica, Fondazione A. Gemelli, Università Cattolica del Sacro Cuore, Rome, Italy

⁵UOC Genetica Medica, Fondazione Policlinico Universitario "A. Gemelli" IRCCS, Rome, Italy

⁶Fetal and Perinatal Pathology Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy

⁷Department of Integrated Surgical and Diagnostic Sciences, University of Genoa, Genoa, Italy

⁸Telethon Institute of Genetics and Medicine (TIGEM), Pozzuoli, Italy

⁹Fetal and Perinatal Pathology Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy

¹⁰Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics and Maternal-Infantile Sciences, University of Genoa, Genoa, Italy

¹¹Department of Vascular Rehabilitation, San Giovanni Battista Hospital, Rome, Italy

¹²Department of Medicine, University of Arizona, Tucson, AZ, USA

¹³Department of Developmental and Social Psychology, Faculty of Medicine and Psychology, Sapienza University of Rome, Rome, Italy

Abstract. – OBJECTIVE: The aim of this qualitative review is to provide an update on the current understanding of the genetic determinants of lipedema and to develop a genetic test to differentiate lipedema from other diagnoses.

MATERIALS AND METHODS: An electronic search was conducted in MEDLINE, PubMed, and Scopus for articles published in English up to March 2019. Lipedema and similar disorders included in the differential diagnosis of lipedema were searched in the clinical synopsis section of OMIM, in GeneCards, Orphanet, and MalaCards.

RESULTS: The search identified several genetic factors related to the onset of lipedema and highlighted the utility of developing genetic diagnostic testing to help differentiate lipedema from other diagnoses.

CONCLUSIONS: No genetic tests or guidelines for molecular diagnosis of lipedema are currently available, despite the fact that genetic testing is fundamental for the differential diagnosis of lipedema against Mendelian genetic obesity, primary lymphedema, and lipodystrophies.

Key Words:

Lipedema, Lipoedema, Differential diagnosis, Subcutaneous fat, Genetic diagnosis.

Introduction

Lipedema is an underdiagnosed chronic debilitating disease characterized by bruising and pain and excess of subcutaneous adipose tissue of the legs and/or arms in women during or after times of hormone change, especially in puberty¹. A definition of lipedema has not yet been included in the 10th revision of the International Classification of Diseases (ICD) by the World Health Organization (WHO) although an ICD-11 code of EF02.2 has been proposed. The first guidelines on lipedema were proposed in 2015 in Germany and by others in 2017 using the international classification of functioning, disability and health²⁻⁴. Lipedema coexists frequently with obesity and can be easily confused with it. However, lipedema is distinguishable from obesity because it is located primarily in the lower limbs and upper extremities sparing the trunk; lipedema minimally responds to diets. Finally, women with lipedema can have a normal Body Mass Index (BMI)⁵. Lipedema can be confused with lymphedema. There are substantial differences between these two diseases:

while lipedema is always bilateral, lymphedema can be unilateral or bilateral; distinctive features of lipedema are pain and bruising that are absent in lymphedema; and in lipedema, the Stemmer sign is negative⁶. Furthermore, lipedema may be associated with multiple lipomas on the arms and/or trunk which are not found in lymphedema^{5,6}. The prevalence of lipedema has been reported to be 1-9/100000⁷. However, it is notable that epidemiological data can fail to recognize lipedema or include individuals who are misdiagnosed with other similar diseases, most commonly obesity, lymphedema, and lipodystrophies. Lipedema can be considered as one component of a spectrum of diseases that are characterized by the dysregulated proliferation of adipose tissue and the presence of pain. The *adiposis dolorosa* (painful fat) spectrum of diseases includes: 1) “Generalized diffuse form of Dercum disease” where painful pearl-sized nodular subcutaneous adipose tissue is widespread throughout the body without larger masses; 2) “generalized nodular form of Dercum disease” where larger painful nodules are found on the arms, trunk, and thighs; 3) “lipedema”, a localized form of painful fat with characteristic distribution of pearl-sized nodular fat and larger masses on the limbs; 4) “localized nodular form of Dercum disease”, usually found around joints (juxta-articular type); and 5) “Madelung disease” or multiple symmetric lipomatosis with nodular fat and lipomas on the upper aspect of the body although a “gynoid type” has been identified^{8,9} (Figure 1). With a variety of diseases of painful fat, diagnosing painful lipedema may be difficult; it would be useful to develop a genetic test that analyses all the genes known for Mendelian non-syndromic genetic obesity, primary lymphedema, and lipodystrophies when trying to diagnose a patient with lipedema, due to the clinical findings these diseases hold in common¹⁰. This review aims to outline current knowledge about the genetics of lipedema and highlights the importance of genetic testing for a more precise diagnosis of lipedema to separate it from similar appearing conditions.

Materials and Methods

This is a “qualitative review” in which, information published about lipedema in referenced scientific journals until March 2019 were collected. An electronic search was conducted in

MEDLINE, PubMed and Scopus, using different combinations of the search terms and concepts “lipedema”, “lipedema genes”, “lipedema etiology”, “lipedema differential diagnosis”, “lipedema and obesity”, “lipedema and lymphedema”, “lipedema and lipodystrophy”, “clinical manifestations of lipedema” and “lipedema genetic testing”. Articles in English were first chosen by examining the title and abstract and, subsequently, analyzed by two independent readings of the whole text. Reference lists were scanned to retrieve other relevant articles. The reviewers assessed the full-text articles. A search for lipedema and similar disorders that overlap in the differential diagnoses was conducted in the clinical synopsis section of OMIM, GeneCards, Orphanet and MalaCards. Orphanet GeneReviews, and Genetic Home Reference were searched to find existing genetic diagnostic tests for lipedema. A general PubMed research was used to investigate clinical, diagnostic and genetic testing guidelines regarding lipedema. Any disagreements were solved through discussion until consensus.

Results

Clinical Aspects of Lipedema

Lipedema is a chronic condition of deposits of painful fat, primarily in the gynoid region, occurring at times of hormone change in women, especially at the time of puberty, that causes gait and joint abnormalities, and which eventually can result in the development of lymphedema, called lipo-lymphedema. Clinically, lipedema has several features: abnormal and painful adipose tissue on the legs, thighs, buttocks, and on the arms in 80%, of cases with a BMI that can be normal or increased without abdominal obesity¹¹; chronic pain that can significantly impact mobility; joint hypermobility associated with fatigue^{1,12,13}; bruising due to increased capillary fragility¹⁴; edema in advanced stages or longstanding lipedema¹⁵ and psychosocial stress and associated psychiatric disorders, especially anxiety and depression¹⁶. The main classifications of lipedema are Schingale classification (Types I, II, III, IV, and V) based on the different distribution of adipose tissue² and Schmeller and Meier-Vollrath classification (Stages I, II, III, and IV) depending on the severity of the disease¹². Conservative treatments of lipedema include: manual lymphatic drainage as part of complex decongestive therapy, subcutaneous adipose tissue therapy to reduce fibrosis and increase




Figure 1. Depiction of phenotypic overlap of different conditions characterized by subcutaneous fat accumulation.

fluid flow through the tissue, compression garment therapy or wrapping to reduce tissue fluid, physical activity and healthy food plans to reduce any obesity component of lipedema, medical foods and supplements (such as selenium, Butcher's broom or diosmin), deep breathing and psychosocial counseling. For patients with minimal or no improvement with conservative treatment, surgical treatments including liposuction and reductive surgery are needed². Lipedema is thought to be either primary (idiopathic, probably genetic) or secondary, associated with hormonal disease, connective tissue disorders and/or autoimmune disease^{17,18}. The long-term effects of lipedema on lifespan are not known¹².

Etiopathogenesis of Lipedema

Lipedema is a chronic disease of lipid metabolism resulting in the symmetrical deposition of subcutaneous adipose tissue associated with hyperplasia and/or hypertrophy of fat cells¹⁹. Etiopathogenesis is unclear but may be associated with impairment of a hormonal axis, since lipedema mainly affects females and its onset is primarily around the time of puberty²⁰. Estrogen has a direct effect on adipose tissue through estrogen receptors (ERs)²¹. Rarely, men are affected with lipedema. Men who develop lipedema tend to be hypogonadal or have liver disease resulting in relatively elevated estrogen levels²². Estrogen may,

therefore, play a role in the development of fat and other tissues that have sex hormone receptors²³. Almost certainly an altered ER pattern or responsiveness exists centrally, not only in the adipose tissue, which may provide an additional explanation for difficulties in gynoid and arm fat loss in patients suffering from lipedema^{10,20}. Additional mechanisms thought to play a role in the pathogenesis of lipedema include increased vascular permeability and damage (microangiopathy), excessive lipid peroxidation, and disturbances in adipocyte metabolism and cytokine production^{20,21}. In a recent study, cells of the stromal vascular fraction (SVF), which includes adipose-derived stromal stem and other cells from 30 patients with lipedema and 22 controls, were characterized. Of note, the yield of stem cells was higher from lipedema fat, which might be explained by an enhanced expression of the mesenchymal (connective tissue) marker CD90 and the endothelial/pericytic marker CD146 suggestive of growth. These markers specifically suggest there may be a chronic capillary injury in lipedema leading to an increased need for repair and protection. Despite these findings, the SVF is thought to be a suitable option for autologous tissue regeneration strategies in lipedema patients²². In summary, lipedema is a heterogeneous likely genetic disease, which manifests in parallel with female hormonal changes and leads to vasculo- and lymphangiopathy. Inflammation of the peripheral nerves and sympathetic innervation abnormalities of the subcutaneous adipose tissue may be responsible for neuropathy with adipocyte hyperproliferation as a secondary phenomenon²⁰.

Genetics of Lipedema

Primary Lipedema

Lipedema is thought to be an inherited disease in most cases. Self-reported positive family history of lipedema is found for up to 64% of women, therefore, a genetic etiology for lipedema is strongly suggested¹⁰. A possible link between the genetic factors and the hormonal role of estrogens in the development of lipedema has been hypothesized²⁴. Autosomal dominant inheritance with incomplete penetrance and sex limitation (most affected female family members) is the most likely mode of inheritance^{12,25}. Lipedema may be differentiated into non-syndromic, syndromic and/or associated with comorbidities. To date, the gene associated with primary isolated lipedema has not been identified. For the diagnosis and research of the genetic basis of lipedema, it is necessary a

multidisciplinary approach that evaluates the genome, the transcriptome, and the methylome.

Syndromic Lipedema

Lipedema may be present within a syndromic context. These diseases may be good models to determine the molecular pathways potentially involved in isolated non-syndromic lipedema.

Example 1: One report associated a germline missense variant, c.196C>T (p.Pro24Leu), in *POU1F1A* in a family with recurrence of short stature and swelling of legs which only affected females in four generations of the family. The only male found affected did not have a lipedema phenotype, only short stature. Affected family members also had growth hormone deficiency, secondary hypothyroidism and hypoprolactinemia²⁶. *POU1F1A* encodes Pit-1, a transcription factor specifically expressed in the anterior pituitary gland, that regulates the expression of the growth hormone (GH), prolactin (PRL) and thyroid-stimulating hormone (TSH) beta-subunit genes^{27,28}.

A pathogenic mutual translocation 46; XX; t(10;15) (q25.2;q11.2) (ClinVar ID: SCV000320871.1) is associated with various conditions, including lipedema²⁹. The 10q25.2 and 15q11.2 bands include several genes that are important for fat metabolism. For instance, changes in the chromosomal region 15q11.2, with loss of gene activity are associated with specific syndromes, such as Prader-Willi syndrome (PWS), characterized by pathologic hyperphagia and obesity³⁰. The genes that map in this region may be good candidates for the onset of lipedema. This is also supported by the observation of one of the authors, that report that lipedema is present in 5% of PWS patients (personal observation).

Example 2: Sotos syndrome 1 is caused by germline mutations in *NSDI*. A novel missense mutation, (p.Cys2175Ser) has been reported in a patient with a familial form of Sotos syndrome characterized by normal intelligence, insulin-dependent diabetes, bronchial asthma, and lipedema. *NSDI* encodes a protein that, together with associated co-regulators, enhances androgen receptor transactivation, therefore it may be partly responsible for the estrogen-mediated lipedema tissue growth³¹.

Example 3: Williams-Beuren syndrome is a multi-systemic neurodevelopmental and metabolic disorder. In some cases, lipedema can be present^{17,32}. This syndrome is caused by deletion of a stretch of 27 genes in the chromosomal region 7q11.23³³ and the best candidates to explain the

involvement of subcutaneous fat tissue are *ELN*³⁴, *FZD9*³⁵, *MLXIPL*³⁶.

Example 4: Pseudoxanthoma elasticum (PXE) is another syndrome in which lipedema can be present. PXE mainly affects skin, eyes, and the cardiovascular and gastrointestinal systems and patients show a progressive mineralization of elastic fibers. Abnormal elastic fibers similar to those found in PXE have been observed in skin lesions of patients with lipedema; furthermore, PXE fibroblasts show an increased cholesterol biosynthesis which could be important in the development of lipedema fat tissue³⁷. PXE can be either autosomal dominant or recessive and is due to germline mutations in *ABCC6* and a subclinical form of PXE has been reported as predisposing to lipedema^{18,38}. Although, to date, no mutations have been reported in *ABCC6*, in association with lipedema.

Example 5: Cutis laxa type III is a multisystem connective tissue disorder characterized by wrinkled, inelastic skin, associated with a neurologic component, retinopathy and the presence of abnormal fat pads on the buttocks and upper thighs. Cutis laxa type III can be either autosomal recessive or dominant and is caused by germline mutations in *ALDH18A1*³⁹, a gene that encodes an enzyme that catalyzes the reduction of glutamate to delta1-pyrroline-5-carboxylate, a critical step in the *de novo* biosynthesis of proline, ornithine, and arginine.

One study reports that hypermobility may be associated with lipedema; in this report, about 50% of women with lipedema had joint hypermobility suggesting that they may also have the common disorder known as hypermobile Ehlers Danlos syndrome (hEDS) formerly hypermobility type or type III^{13,21}. Germline truncating mutations in *TNXB* and a single germline variant in *COL3A1*, (p.Gly637Ser) have been found to cause the Ehlers-Danlos syndrome type III, to date, no mutations have been found in those two genes in association with lipedema^{40,41}.

Clinically Overlapping Conditions with Lipedema

In thinking about a possible genetic test for lipedema, it is important to consider the genes mutated in conditions that may be phenotypically overlapping or confused with lipedema for differential diagnosis, such as lipomas or familial multiple lipomatosis.

For instance, disruption of *HMG2*, located in the chromosomal band 12q14.3, has frequently

been detected in human benign tumors of mesenchymal origin, including lipomas^{42,43}.

A rare case of autosomal dominant familial multiple subcutaneous lipomatosis (FML) and increased predilection to cancers co-segregating with a novel germline truncating mutation in *PALB2* (c.2716delT) has been reported⁴⁴.

Two patients with encephalocraniocutaneous lipomatosis, a congenital hamartomatous disorder characterized by unilateral skin lesions, lipomas, and ipsilateral ophthalmological and cerebral malformations, have been found to carry a *KRAS* somatic mutation c.436G>A (p.Ala146Thr) and a *NF1 de novo* germline truncating mutation, respectively⁴⁴⁻⁴⁷.

PIK3CA somatic activating mutations, previously reported in cancer and overgrowth syndromes, have been discovered in the affected tissue of individuals with facial infiltrating lipomatosis⁴⁸. The syndrome of multiple symmetric lipomatosis, partial lipodystrophy and insulin resistance is caused by germline mutations in *LIPE*. These mutations result in marked inhibition of lipolysis from adipose tissue depots⁴⁹ which could explain fat expansion in lipedema.

The Pierpont syndrome, a plantar lipomatosis with limb abnormalities, unusual facies and developmental delay is caused by a *de novo* heterozygous missense mutation (p.Tyr446Cys) in *TBLIXR1*⁵⁰. In two cases, two different mutations have been reported p.Cys325Tyr and p.Tyr446His⁵¹.

Hereditary retinoblastoma is caused by variants in *RBI* (40% of mutations are germline, inherited or *de novo*; 60% of mutations are somatic). Retinoblastoma, an embryonic malignant neoplasm of retinal origin, is found in association with lipomatosis in less than 4% of the affected subjects⁵². Lipomatosis predisposition in hereditary retinoblastoma is not associated with specific *RBI* variants but is probably determined by modifying factors⁵³. Interestingly, the retinoblastoma interacting zinc finger gene *RIZ* codes for a protein that is a downstream effector of estrogen action and is involved in an alternative intracellular pathway, mediating the mitogenic effect of estradiol^{54,55}, a finding that may be pertinent to lipedema.

In support of the hypothesis that multiple symmetric lipomatosis is caused by mitochondrial mutations^{10,56}, there are several studies. A study reported that 32.4% (11/34) of cases affected by myoclonic epilepsy with ragged-red fibers (MERF) show the presence of multiple lipomatosis.

Of them, only 6 had isolated lipomatosis with no myopathic complaints and/or CNS involvement. All of these patients carried the m.8344A>G mutation in the MTTK, transcribed in the tRNA^{Lys57-59}. In a multigenerational Canadian family with 8 affected members, the complete co-segregation of the rare mitochondrial variant in MTTK (8344A>G) and the phenotype was reported⁶⁰.

Multiple symmetric lipomatosis (Madelung disease), a rare disorder of adipocyte differentiation characterized by benign, diffuse lipomatosis in the cephalic, cervical, and upper thoracic subcutaneous depots, can carry biallelic mutations in MFN2, a gene encoding a mitochondrial protein already associated with Charcot-Marie-Tooth syndrome^{45,61}.

Familial Partial Lipodystrophies and Lipedema

It is important to point out that in some cases of partial lipodystrophy there may be an accumulation of subcutaneous fat tissue in specific depots coupled with the reduction of fat in other regions. The majority of familial partial lipodystrophy (FPLD) syndromes are autosomal dominant. Phenotypically, patients with FPLD, type 2 and some of the other partial lipodystrophies lack limb and gluteal subcutaneous fat, while in FPLD, type 1, the abdomen shows weight gain and the limbs and buttocks lose fat⁶². In most cases, body fat redistribution occurs during puberty. Adipose tissue in the face, neck and intra-abdominal areas may be increased in FPLD, type 2. Diabetes and metabolic complications, such as coronary artery disease, occur in adulthood in these disorders⁶³.

FPLD, type 2 is caused by autosomal dominant mutations in LMNA, the most common being p.Arg482Gln; this gene encodes the nuclear lamin A/C protein. Fat loss typically occurs in the buttocks and limbs, and gain of fat occurs to the face, neck, abdominal viscera, and labia, although there is phenotypic variability. Patients may also have lipomas⁶³. While lipedema may be misdiagnosed as lipodystrophy, the metabolic differences between lipedema, which confers low cardiovascular risk, and FPLD, type 2, which confers high cardiovascular risk, are clear.

FPLD, type 3 is caused by heterozygous missense mutations in PPAR γ . Fat loss usually occurs with loss of fat from the trunk, buttocks, and limbs and there is no increase in head and neck fat. However, there may be a gain of fat on the abdomen^{63,64}. The phenotype of women with FPLD3 is different from that of women with lipedema,

with upper body obesity compared to gynoid fat deposits, respectively.

FPLD4 is caused by heterozygous truncating mutations in PLIN1, encoding perilipin-1. This protein covers lipid storage droplets in adipocytes, protecting them until they can be broken down by hormone-sensitive lipases. Perilipin-1 is the major cAMP-dependent protein kinase substrate in adipocytes and may play a role in the inhibition of lipolysis⁶³. The phenotype of women with FPLD4 is different from that of women with lipedema, with upper body obesity compared to gynoid fat deposits, respectively.

FPLD5 has been reported in a single case caused by a homozygous truncating mutation in CIDEA (p.Glu186*). Perilipin-1 and CIDEA are involved in the structure and function of adipocyte lipid droplets⁶³.

FPLD6 is caused by homozygous truncating mutations in LIPE, encoding hormone-sensitive lipase, which is involved in the regulation of lipolysis⁶³.

Akt2-linked lipodystrophy was described in a single family and is caused by heterozygous missense mutations in AKT2, which is a key mediator of insulin signaling downstream of the insulin receptor⁶⁵. The phenotype of women with FPLD6 is different from that of women with lipedema, with upper body obesity compared to gynoid fat deposits, respectively.

Familial partial lipodystrophies share common phenotypes. The typical loss of fat occurs to the buttocks and limbs and a gain of fat to the face, neck, and abdomen. Metabolic complications of the familial partial lipodystrophies include insulin resistance, diabetes mellitus, hypertriglyceridemia, hepatic steatosis, hypertension, and coronary artery disease, none of which correlates with the phenotype of lipedema. However, genes important in partial lipodystrophy may be important in modifying the phenotype of lipedema.

In an African-American pedigree with a novel autosomal dominant atypical FPLD, Garg et al⁶⁶ performed linkage analysis for candidate regions and whole-exome sequencing to identify the disease-causing mutation. The affected individuals had marked loss of the fat from the extremities, with excess fat in the face and neck, and metabolic complications. The authors identified a mutation (c.202C>T, p.Leu68Phe) in the adrenoceptor α 2A (ADRA2A) in all affected subjects but not in the unaffected. ADRA2A regulates norepinephrine release and its activation reduces lipolysis in adipocytes. The mutation p.Leu68Phe presumably

causes a rare atypical FPLD by inducing excessive lipolysis in certain adipose tissue depots⁶⁶.

Candidate Genes from Genome-Wide Association and Animal Models Studies

Several types of studies have been conducted to identify candidate genes associated with lipedema: genome-wide association studies (GWAS) and studies on animal models.

LYPLALI, *TBX15*, *HOXC13*, *RSPO3*, *CPEB4*, *VEGFA*, *STAB1*, *GRB14*, *ZNRF3*, and *PIGC* have been associated with subcutaneous adipose depots by genome-wide association studies and confirmed by functional analyses. The association of these genes with body fat distribution is independent of any effects of BMI and/or obesity, demonstrating genetic control of body fat distribution, distinct from that of overall adiposity^{67,68}.

ATXN1 and *UBE2E2* play a physiological role in adipogenesis. In support of these genes as candidate genes for lipedema, seven *loci* associated with ectopic fat traits have been identified in a multi-ethnic, sample-size-weighted, fixed-effects genome-wide association meta-analysis study at the genomic level: *ATXN1*, *UBE2E2*, *EBF1*, *RREB1*, *GSDMB*, *GRAMD3*, *ENSA*. Functional analyses of *ATXN1* and *UBE2E* showed that loss of *Atnx1* or *Ube2e2* in primary mouse adipose progenitor cells hindered differentiation of adipocytes⁶⁹.

VEGFR3 signaling is fundamental for the development and maintenance of the lymphatic vascular system⁷⁰. *VEGFR3* is associated with the development of lymphedema in humans⁷¹. Mice with *VEGFR-3* gene heterozygous inactivating missense mutations have hypoplastic dermal lymphatics and thickening of the subcutaneous adipose tissue. Vascular endothelial growth factors *VEGFC* and *VEGFD* stimulate lymphangiogenesis, whereas *VEGFR3* is a potent inhibitor of *VEGFC/VEGFD* signaling. In the skin of transgenic mice, *VEGFR3* inhibits fetal lymphangiogenesis and induces regression of already formed lymphatic vessels, though the blood vasculature remains normal. The transgenic K14-*VEGFR3*-Ig mice show, besides other features, increased deposition of subcutaneous fat⁷⁰.

The ablation of *PRDM16* in mice adipocytes has significant reductions in thermogenic gene expression and O₂ consumption of white adipose tissue, both in the basal state and following stimulation with cold and β 3-adrenergic agonist treatment resulting in marked enlargement of the subcutaneous adipose tissue. These animals

developed obesity on a high-fat diet, with severe insulin resistance and hepatic steatosis, which is, however, not seen in lipedema in the absence of obesity. However, they also showed altered fat distribution⁷².

Fat-specific *Sirt6* knockout mice are sensitized to high-fat diet-induced obesity resulting in adipocyte hypertrophy caused by compromised lipolytic activity due to the decreased expression of the lipolytic enzyme, adipose tissue triglyceride lipase (*ATGL*). Fat-specific *Sirt6* KO also showed increased inflammation in adipose tissue, which contributes to insulin resistance in high-fat diet-fed mice. These findings are consistent with the reduced expression of *SIRT6* and *ATGL* observed in obese patients⁷³. Recently, in humans, it has been observed that, in the subcutaneous adipose tissue, increased body mass index modifies the expression of the genes encoding sirtuins and their target genes, which are metabolic regulators of adipose tissue⁷⁴. Interestingly, *SIRT6* is widely expressed in retinoblastomas, a tumor that in 4% of cases shows the presence of lipomatosis.

FGF21 improves systemic insulin sensitivity by promoting the healthy expansion of subcutaneous adipose tissue. *FGF21* knockout mice show less subcutaneous adipose tissue and are more insulin-resistant when fed with a high-fat diet. Moreover, serum *FGF21* levels positively correlate with the subcutaneous adipose tissue area in insulin-sensitive obese individuals⁷⁵.

IL18 receptor 1- knockout mice have an increased susceptibility to dietary-induced obesity. It was observed by Pazos et al⁷⁶ that *Il18* knockout mice were extremely dietary obesity-prone and did not develop diet-induced thermogenesis as assessed by brown adipose tissue and white adipose tissue *Ucp1* mRNA levels. Interestingly, women with lipedema have reported cold areas in fat tissue.

PANX1 encodes a glycoprotein, pannexin 1, which plays a key role in regulating the formation of adipocytes and fat accumulation. Increased adipocyte size and decreased adipocyte numbers were observed in subcutaneous fat of the *Panx1* knockout mice compared to wild-type mice. Knockout mice lacking *Panx1* gene have significantly greater total fat mass and reduced lean mass under a normal diet. By metabolic cage data, it was observed that *Panx1* knockout mice on a high-fat diet display significantly increased activity levels, higher ambulatory activity, and reduced sleep duration with respect to wild type mice. Absence of the *Panx1* protein also leads to increased insulin and blood glucose levels, which increase the risk of type 2 diabetes⁷⁷.

SCD1 encodes an enzyme that catalyzes the synthesis of monounsaturated fatty acids and is an important regulator of whole-body energy homeostasis. Knockout mice lacking *Scd1* gene display marked sebaceous gland hypoplasia, depletion of sebaceous lipids, increased energy expenditure and protection from high-fat diet-induced obesity, and severe cold intolerance. mRNA levels of *SCD1* in subcutaneous adipose tissue correlate with BMI in morbidly obese subjects⁷⁸. *SCD1* mRNA levels have not been tested in lipedema but loss of *SCD1* activity was inferred due to the lower levels of unsaturated fatty acids from subcutaneous adipose tissue of people with fat disorders⁷⁹.

PROX1 is a gene correlated to leaky lymphatic vessels as observed by studying mice with functional inactivation of a single allele of this homeobox gene. *PROX1* is fundamental for lymphatic vascular development and *Prox1* knockout mice are obese. *Prox1* haploinsufficiency results in subcutaneous and intra-abdominal fat accumulation in mice⁸⁰, a potential subclinical cause for the accumulation of lipedema fat tissue.

ZFP423 encodes a multi-zinc-finger transcription factor expressed in preadipocytes and mature adipocytes *in vivo* and is responsible for the terminal differentiation of subcutaneous white adipose tissue. Knocking out *Zfp423* in mice results in white-to-beige phenotypic adipocytes switch. When *Zfp423* is removed from precursors and/or actively differentiating cells, there is a defect in the terminal differentiation of white adipocytes. The loss of lipid in the adipocytes in the absence of *Zfp423* is due to lower expression of *PPAR γ* and other adipocyte-selective genes. Therefore, *ZFP423* has a fundamental role in the establishment and maintenance of the white adipose lineage⁸¹ and could be speculated to play a role in maintenance of lipedema fat.

Lipedema Diagnosis and Genetic Test

The diagnosis of lipedema is mainly based on the medical history and physical examination of the patient². Multiple imaging techniques such as ultrasound, magnetic resonance imaging (MRI) that allows identification of subclinical nodules, computed tomography, lymphoangioscintigraphy, indirect lymphangiography, and dual energy X-ray absorptiometry have all been applied in the diagnosis and study of lipedema. Fluorescence micrography makes it possible to detect multiple microaneurysms in lymphatic capillaries in the skin regions affected by lipedema not detectable in other pathologies, such as scleroderma and ve-

nous insufficiency⁸². In some cases, the MRI analysis of excised lipomatosis nodules revealed that they were angioliipomas identifiable with MRI⁸³. In another work, using MRI in women with lipedema, higher levels of sodium were found in the fat tissue with even early stage lipedema compared to controls⁸⁴ suggesting a strong role for lymphatic dysfunction in lipedema⁸⁵.

The Streeten test may be considered as a functional test to differentiate lipedema from lymphedema². General blood tests are important because patients with lipedema may have diseases that are responsible for the worsening of their symptoms².

Characterization of family history and clinical phenotype will be fundamental for genetic testing in lipedema, as a causal mutation is thought to be important in this disease²⁵. Currently, genetic tests and guidelines for the molecular diagnosis of lipedema are not available. However, genetic testing to exclude overlapping conditions or syndromic lipedema would be of fundamental importance. Lipedema is often misdiagnosed because of its similarity with primary lymphedema, Mendelian non-syndromic obesity, and generalized and partial lipodystrophies. We propose the use of a genetic test based on the next generation sequencing of a panel of genes that could be involved in syndromic lipedema and are involved in phenotypically overlapping disorders (obesity, lymphedema, and lipodystrophies) with similar adiposity to exclude these pathologies. After the exclusion of the presence of known pathogenic mutations in both blood (germline mutations) and affected tissue (somatic mutations), whole exome/genome sequencing would be required to identify the still-unknown gene (2). In Figure 2, a possible and ideal diagnostic genetic test for lipedema is schematized.

Conclusions

Lipedema is a very disabling disease that affects the quality of life, but unfortunately, it may be easily misdiagnosed¹. It belongs to a spectrum of the localized painful depots of subcutaneous adipose tissue (*adiposis dolorosa*) that includes a generalized/localized/diffuse/nodular form of Dercum disease, Madelung disease (multiple symmetric lipomatosis), partial lipodystrophy, familial multiple lipomatosis, and lymphedema. A timely and accurate diagnosis of lipedema is essential for early intervention and treatment of the disease to prevent the development

of co-morbidities. For this purpose, it is important to develop a genetic test for lipedema based on increasing knowledge of causative genes for other disorders and rapid advances in genetic screening technologies. Lipedema is an inherited disease expressed mainly in females, with males serving as carriers^{27,86}. Lipedema is a relatively poorly studied disease, in fact, the mechanisms concerning the etiopathogenesis and the genetics of lipedema are not yet fully

known and there are no current reliable biomarkers or imaging modalities. Currently, the gene/s associated with primary lipedema are not known, whereas some genes associated with syndromic lipedema are known. Genome-wide association and animal models studies have made it possible to identify possible candidate genes and genetic pathways involved in the onset of lipedema. The advent of new molecular technologies will improve research and diagnostic




Figure 2. Scheme of a possible and ideal diagnostic genetic test for genetic subcutaneous fat accumulation.

process, ensuring the possibility of more effective prevention and treatment with clear improvements in the management of lipedema patients. The development of a genetic test containing candidate genes for lipedema and causative genes of diseases that can be confused with lipedema will be fundamental to find the etiology of lipedema and to provide patients with more targeted care and follow-up. By analysis of the whole exome with NGS technology in the patient and their parents could allow identification of new candidate genes for lipedema⁸⁷ but the phenotype must be clearly defined. A multidisciplinary approach and with different technologies would be needed. Understanding the molecular mechanisms underlying the onset of lipedema requires a multi-pronged approach based on, for instance, genome, transcriptome and methylome studies. Hopefully, the use of NGS technologies will help clinicians classify people that meet the criteria for the *adipositas dolorosa* spectrum.

Conflict of Interests

The Authors declare that there are no conflicts of interest.

References

- 1) WOLD LE, HINES EA JR, ALLEN EV. Lipedema of the legs; a syndrome characterized by fat legs and edema. *Ann Intern Med* 1951; 34: 1243-1250.
- 2) HALK AB, DAMSTRA RJ. First Dutch guidelines on lipedema using the international classification of functioning, disability and health. *Phlebology* 2017; 32: 152-159.
- 3) REICH-SCHUPKE S, SCHMELLER W, BRAUER WJ, CORNELI ME, FAERBER G, LUDWIG M, LULAY G, MILLER A, RAPPRICH S, RICHTER DF, SCHACHT V, SCHRADER K, STÜCKER M, URE C. S1 guidelines: lipedema. *J Dtsch Dermatol Ges* 2017; 15: 758-767.
- 4) HARDY D, WILLIAMS A. Best practice guidelines for the management of lipoedema. *Br J Community Nurs* 2017; 22: S44-S48.
- 5) WARREN AG, JANZ BA, BORUD LJ, SLAVIN SA. Evaluation and management of the fat leg syndrome. *Plast Reconstr Surg* 2007; 119: 9e-15e.
- 6) FORNER-CORDERO I, SZOLNOKY G, FORNER-CORDERO A, KEMÉNY L. Lipedema: an overview of its clinical manifestations, diagnosis and treatment of the disproportional fatty deposition syndrome - systematic review. *Clin Obes* 2012; 2: 86-95.
- 7) ORPHANET. Lipedema. [https://www.orpha.net/consor/cgi-bin/Disease_Search.php?lng=EN&data_id=11100&Disease_Disease_Search_diseaseGroup=Lipoedema&Disease_Disease_Search_diseaseType=Pat&Disease\(s\)/group%20of%20diseases=Lipedema&title=Lipedema&search=Disease_Search_Simple](https://www.orpha.net/consor/cgi-bin/Disease_Search.php?lng=EN&data_id=11100&Disease_Disease_Search_diseaseGroup=Lipoedema&Disease_Disease_Search_diseaseType=Pat&Disease(s)/group%20of%20diseases=Lipedema&title=Lipedema&search=Disease_Search_Simple) [Accessed 21 March 2019].
- 8) HANSSON E, SVENSSON H, BRORSON H. Review of Dercum's disease and proposal of diagnostic criteria, diagnostic methods, classification and management. *Orphanet J Rare Dis* 2012; 7: 23.
- 9) BELTRAN K, HERBST KL. Differentiating lipedema and Dercum's disease. *Int J Obes* 2017; 41: 240-245.
- 10) HERBST KL. Rare adipose disorders (RADs) masquerading as obesity. *Acta Pharmacol Sin* 2012; 33: 155-172.
- 11) DIETZEL R, REISSHAUER A, JAHR S, CALAFIORE D, ARMBRECHT G. Body composition in lipoedema of the legs using dual-energy X-ray absorptiometry: a case-control study. *Br J Dermatol* 2015; 173: 594-596.
- 12) SCHMELLER W, MEIER-VOLLRATH I. *Lymphedema. Diagnosis and therapy.* Viavital Verlag GmbH, 2007.
- 13) TORRE YS, WADEEA R, ROSAS V, HERBST KL. Lipedema: friend and foe. *Horm Mol Biol Clin Investig* 2018; 33.pii:/hmbci.2018.33.issue-1/hmbci-2017-0076/hmbci-2017-0076.xml. doi: 10.1515/hmbci-2017-0076.
- 14) WOLLINA U. Lipedema-an update. *Dermatol Ther* 2019; 32: e12805.
- 15) RAPPRICH S, DINGLER A, PODDA M. Liposuction is an effective treatment for lipedema results of a study with 25 patients. *J Dtsch Dermatol Ges* 2011; 9: 33-40.
- 16) DUDEK JE, BIAŁASZEK W, OSTASZEWSKI P. Quality of life in women with lipoedema: a contextual behavioral approach. *Qual Life Res* 2016; 25: 401-408.
- 17) WAXLER JL, GUARDINO C, FEINN RS, LEE H, POBER BR, STANLEY TL. Altered body composition, lipedema, and decreased bone density in individuals with Williams Syndrome: a preliminary report. *Eur J Med Genet* 2017; 60: 250-256.
- 18) TAYLOR NE, FOSTER WC, WICK MR, PATTERSON JW. Tumefactive lipedema with pseudoxanthoma elasticum-like microscopic changes. *J Cutan Pathol* 2004; 31: 205-209.
- 19) SZÉL E, KEMÉNY L, GROMA G, SZOLNOKY G. Pathophysiological dilemmas of lipedema. *Med Hypotheses* 2014; 83: 599-606.
- 20) MAYES JS, WATSON GH. Direct effects of sex steroid hormones on adipose tissues and obesity. *Obes Rev* 2004; 5: 197-216.
- 21) HERBST KL, MIRKOVSKAYA L, BHARHAGAVA A, CHAVA Y, CHT TE. Lipedema fat and signs and symptoms of illness, increase with advancing stage. *Arch Med* 2015; 7: 10.
- 22) FIFE CE, MAUS EA, CARTER MJ. Lipedema: a frequently misdiagnosed and misunderstood fatty deposition syndrome. *Adv Skin Wound Care* 2010; 23: 81-92.
- 23) SHIN BW, SIM YJ, JEONG HJ, KIM GC. Lipedema, a rare disease. *Ann Rehabil Med* 2011; 35: 922-927.
- 24) PRIGLINGER E, WURZER C, STEFFENHAGEN C, MAIER J, HOFER V, PETERBAUER A, NUERNBERGER S, REDL H, WOLBANK S, SANDHOFER M. The adipose tissue-derived stromal vascular fraction cells from lipedema patients: are they different? *Cytotherapy* 2017; 19: 849-860.
- 25) CHILD AH, GORDON KD, SHARPE P, BRICE G, OSTERGAARD P, JEFFERY S, MORTIMER PS. Lipedema: an inherited condition. *Am J Med Genet A* 2010; 152: 970-976.

- 26) BANO G, MANSOUR S, BRICE G, OSTERGAARD P, MORTIMER PS, JEFFERY S, NUSSEY S. Pit-1 mutation and lipodema in a family. *Exp Clin Endocrinol Diabetes*. 2010; 118: 377-380.
- 27) FÖLDI M, FÖLDI E. Földi's textbook of lymphology. Urban & Fischer, 2012.
- 28) GONZÁLEZ-PARRA S, CHOWEN JA, GARCÍA-SEGURA LM, ARGENTE J. In vivo and in vitro regulation of pituitary transcription factor-1 (Pit-1) by changes in the hormone environment. *Neuroendocrinology* 1996; 63: 3-15.
- 29) REDIN C, BRAND H, COLLINS RL, KAMMIN T, MITCHELL E, HODGE JC, HANSCOM C, PILLALAMARRI V, SEABRA CM, ABBOTT M, ABDUL-RAHMAN OA, ABERG E, ADLEY R, ALCARAZ-ESTRADA SL, ALKURAYA FS, AN Y, ANDERSON MA, ANTOLIK C, ANYANE-YEBOA K, ATKIN JF, BARTELL T, BERNSTEIN JA, BEYER E, BLUMENTHAL I, BONGERS EM, BRILSTRA EH, BROWN CW, BRÜGGENWIRTH HT, CALLEWAERT B, CHIANG C, CORNING K, COX H, CUPPEN E, CURRALL BB, CUSHING T, DAVID D, DEARDORFF MA, DHEEDENE A, D'HOOGHE M, DE VRIES BB, EARL DL, FERGUSON HL, FISHER H, FITZPATRICK DR, GERROL P, GIACHINO D, GLESSNER JT, GLIEM T, GRADY M, GRAHAM BH, GRIFFIS C, GRIPP KW, GROPMAN AL, HANSON-KAHN A, HARRIS DJ, HAYDEN MA, HILL R, HOCHSTENBACH R, HOFFMAN JD, HOPKIN RJ, HUBSHMAN MW, INNES AM, IRONS M, IRVING M, JACOBSEN JC, JANSSENS S, JEWETT T, JOHNSON JP, JONGMANS MC, KAHLER SG, KOOLEN DA, KORZELIUS J, KROISEL PM, LACASSIE Y, LAWLESS W, LEMYRE E, LEPPIG K, LEVIN AV, LI H, LI H, LIAO EC, LIM C, LOSE EJ, LUCENTE D, MACERA MJ, MANAVALAN P, MANDRILE G, MARCELIS CL, MARGOLIN L, MASON T, MASSER-FRYE D, MCCLELLAN MW, MENDOZA CJ, MENTEN B, MIDDELKAMP S, MIKAMI LR, MOE E, MOHAMMED S, MONONEN T, MORTENSON ME, MOYA G, NIEUWINT AW, ORDULU Z, PARKASH S, PAUKER SP, PEREIRA S, PERRIN D, PHELAN K, AGUILAR RE, PODDIGHE PJ, PREGNO G, RASKIN S, REIS L, RHEAD W, RITA D, RENKENS I, ROELENIS F, RULIERA J, RUMP P, SCHILIT SL, SHAHEEN R, SPARKES R, SPIEGEL E, STEVENS B, STONE MR, TAGOE J, THAKURIA JV, VAN BON BW, VAN DE KAMP J, VAN DER BURGT I, VAN ESSEN T, VAN RAVENSWAALI-ARTS CM, VAN ROOSMALEN MJ, VERGULT S, VOLKER-TOUW CM, WARBURTON DP, WATERMAN MJ, WILEY S, WILSON A, YERENA-DE VEGA MC, ZORI RT, LEVY B, BRUNNER HG, DE LEEUW N, KLOOSTERMAN WP, THORLAND EC, MORTON CC, GUSSELLA JF, TALKOWSKI ME. The genomic landscape of balanced cytogenetic abnormalities associated with human congenital anomalies. *Nat Genet* 2017; 49: 36-45.
- 30) BUITING K. Prader-Willi syndrome and Angelman syndrome. *Am J Med Genet C Semin Med Genet* 2010; 154: 365-376.
- 31) ZECHNER U, KOHLSCHMIDT N, KEMPF O, GEBAUER K, HAUG K, ENGELS H, HAAF T, BARTSCH O. Familial Sotos syndrome caused by a novel missense mutation, C2175S, in NSD1 and associated with normal intelligence, insulin dependent diabetes, bronchial asthma, and lipedema. *Eur J Med Genet* 2009; 52: 306-310.
- 32) SHAIKH S, WAXLER JL, LEE H, GRINKE K, GARRY J, POBER BR, STANLEY TL. Glucose and lipid metabolism, bone density, and body composition in individuals with Williams syndrome. *Clin Endocrinol (Oxf)* 2018; 89: 596-604.
- 33) MORRIS CA. Williams Syndrome. *GeneReviews*. Seattle (WA): University of Washington, 2017.
- 34) LIU Y, ARON-WISNEWSKY J, MARCELIN G, GENSER L, LE NAOUR G, TORCIVIA A, BALUVOIS B, BOUCHET S, PELLOUX V, SASSO M, MIETTE V, TORDJMAN J, CLÉMENT K. Accumulation and changes in composition of collagens in subcutaneous adipose tissue after bariatric surgery. *J Clin Endocrinol Metab* 2016; 101: 293-304.
- 35) WADA N, HASHINAGA T, OTABE S, YUAN X, KURITA Y, KAKINO S, OHOKI T, NAKAYAMA H, FUKUTANI T, TAJIRI Y, YAMADA K. Selective modulation of Wnt ligands and their receptors in adipose tissue by chronic hyperadiponecตินemia. *PLoS One* 2013; 8: e67712.
- 36) SONG Z, XIAOLI AM, YANG F. Regulation and metabolic significance of de novo lipogenesis in adipose tissues. *Nutrients* 2018; 10(10). pii: E1383.
- 37) KUZAJ P, KUHN J, DABISCH-RUTHE M, FAUST I, GÖTTING C, KNABBE C, HENDIG D. ABCG6- a new player in cellular cholesterol and lipoprotein metabolism? *Lipids Health Dis* 2014; 13: 118.
- 38) TERRY SF, BERCOVITCH L. Pseudoxanthoma elasticum. *GeneReviews*. Seattle (WA): University of Washington, 2012.
- 39) WOLTHUIS DF, VAN ASBECK E, MOHAMED M, GARDEITCHIK T, LIM-MELIA ER, WEVERS RA, MORAVA E. Cutis laxa, fat pads and retinopathy due to ALDH18A1 mutation and review of the literature. *Eur J Paediatr Neurol* 2014; 18: 511-515.
- 40) MALFAIT F, FRANCOMANO C, BYERS P, BELMONT J, BERGLUND B, BLACK J, BLOOM L, BOWEN JM, BRADY AF, BURROWS NP, CASTORI M, COHEN H, COLOMBI M, DEMIRDAS S, DE BACKER J, DE PAEPE A, FOURNEL-GIGLEUX S, FRANK M, GHALI N, GIUNTA C, GRAHAME R, HAKIM A, JEUNEMAITRE X, JOHNSON D, JUUL-KRISTENSEN B, KAPFERER-SEEBACHER I, KAZKAZ H, KOSHO T, LAVALLEE ME, LEVY H, MENDOZA-LONDONO R, PEPIN M, POPE FM, REINSTEIN E, ROBERT L, ROHRBACH M, SANDERS L, SOBEY GJ, VAN DAMME T, VANDERSTEEN A, VAN MOURIK C, VOERMANS N, WHEELDON N, ZSCHOCKE J, TINKLE B. The 2017 international classification of the Ehlers-Danlos syndromes. *Am J Med Genet C Semin Med Genet* 2017; 175: 8-26.
- 41) NARCISI P, RICHARDS AJ, FERGUSON SD, POPE FM. A family with Ehlers-Danlos syndrome type III/articular hypermobility syndrome has a glycine 637-to-serine substitution in type III collagen. *Hum Mol Genet* 1994; 3: 1617-1620.
- 42) BATTISTA S, FIDANZA V, FEDELE M, KLEIN-SZANTO AJ, OUTWATER E, BRUNNER H, SANTORO M, CROCE CM, FUSCO A. The expression of a truncated HMGI-C gene induces gigantism associated with lipomatosis. *Cancer Res* 1999; 59: 4793-4797.
- 43) ASHAR HR, FEJZO MS, TKACHENKO A, ZHOU X, FLETCHER JA, WEREMOWICZ S, MORTON CC, CHADA K. Disruption of the architectural factor HMGI-C: DNA-binding AT hook motifs fused in lipomas to distinct transcriptional regulatory domains. *Cell* 1995; 82: 57-65.
- 44) REDDY N, MALIPATIL B, KUMAR S. A rare case of familial multiple subcutaneous lipomatosis with novel PALB2 mutation and increased predilection to cancers. *Hematol Oncol Stem Cell Ther* 2016; 9: 154-156.
- 45) SAWYER SL, CHEUK-HIM NG A, INNES AM, WAGNER JD, DYMENT DA, TETREAULT M; CARE RARE CANADA CONSORTIUM, MAJEWSKI J, BOYCOTT KM, SCREATION RA, NICHOLSON G. Homozygous mutations in MFN2 cause multiple symmetric lipomatosis associated with neuropathy. *Hum Mol Genet* 2015; 24: 5109-5114.

- 46) LEGIUS E, WU R, EYSSSEN M, MARYNYEN P, FRYNS JP, CASIMAN JJ. Encephalocraniocutaneous lipomatosis with a mutation in the NF1 gene. *J Med Genet* 1995; 32: 316-319.
- 47) BOPPUDI S, BÖGERSHAUSEN N, HOVE HB, PERCIN EF, ASLAN D, DVORSKY R, KAYHAN G, LI Y, CURSIEFEN C, TANTCHEVA-POOR I, TOFT PB, BARTSCH O, LISSEWSKI C, WIELAND I, JAKUBICZKA S, WOLLNIK B, AHMADIAN MR, HEINDL LM, ZENKER M. Specific mosaic KRAS mutations affecting codon 146 cause oculoectodermal syndrome and encephalocraniocutaneous lipomatosis. *Clin Genet* 2016; 90: 334-342.
- 48) MACLELLAN RA, LUKS VL, VIVERO MP, MULLIKEN JB, ZURAKOWSKI D, PADWA BL, WARMAN ML, GREENE AK, KUREK KC. PIK3CA activating mutations in facial infiltrating lipomatosis. *Plast Reconstr Surg* 2014; 133: 12e-9e.
- 49) ZOLOTOV S, XING C, MAHAMID R, SHALATA A, SHEIKH-AHMAD M, GARG A. Homozygous LIPE mutation in siblings with multiple symmetric lipomatosis, partial lipodystrophy, and myopathy. *Am J Med Genet A* 2017; 173: 190-194.
- 50) HEINEN CA, JONGEJAN A, WATSON PJ, REDEKER B, BOELLEN A, BOUDZOVITCH-SUROVITSEVA O, FORZANO F, HORDIJK R, KELLEY R, OLNEY AH, PIERPONT ME, SCHAEFER GB, STEWART F, VAN TROTSENBURG AS, FLIERS E, SCHWABE JW, HENNEKAM RC. A specific mutation in TBL1XR1 causes Pierpont syndrome. *J Med Genet* 2016; 53: 330-337.
- 51) LEMATRE C, THEVENON J, DUFFOURD Y, NAMBOT S, HAQUET E, VUADALLE B, GENEVIEVE D, SARDA P, BRUEL AL, KUENTZ P, WELLS CF, FAIVRE L, WILLEMS M. TBL1XR1 mutations in Pierpont syndrome are not restricted to the recurrent p.Tyr446Cys mutation. *Am J Med Genet A* 2018; 176: 2813-2818.
- 52) TENNA S, BARONE M, GIGLIOFIORITO P, PERSICETTI P. Retinoblastoma plus lipomatosis: an autosomal dominant syndrome. *Arch Plast Surg* 2014; 41: 785-787.
- 53) GENUARDI M, KLUTZ M, DEVRIENDT K, CARUSO D, STIRPE M, LOHMANN DR. Multiple lipomas linked to an RB1 gene mutation in a large pedigree with low penetrance retinoblastoma. *Eur J Hum Genet* 2001; 9: 690-694.
- 54) LIU L, SHAO G, STEELE-PERKINS G, HUANG S. The retinoblastoma interacting zinc finger gene RIZ produces a PR domain-lacking product through an internal promoter. *J Biol Chem* 1997; 272: 2984-2991.
- 55) ABBONDANZA C, MEDICI N, NIGRO V, ROSSI V, GALLO L, PILUSO G, BELSITO A, ROSCIGNO A, BONTEMPO P, PUCA AA, MOLINARI AM, MONCHARMONT B, PUCA GA. The retinoblastoma-interacting zinc-finger protein RIZ is a downstream effector of estrogen action. *Proc Natl Acad Sci U S A* 2000; 97: 3130-3135.
- 56) KLOPSTOCK T, NAUMANN M, SCHALKE B, BISCHOF F, SEIBEL P, KOTTLORS M, ECKERT P, REINERS K, TOYKA KV, REICHMANN H. Multiple symmetric lipomatosis: abnormalities in complex IV and multiple deletions in mitochondrial DNA. *Neurology* 1994; 44: 862-866.
- 57) CHONG PS, VUCIC S, HEDLEY-WHYTE ET, DREYER M, CROS D. Multiple symmetric lipomatosis (Madelung's disease) caused by the MERRF (A8344G) mutation: a report of two cases and review of the literature. *J Clin Neuromuscul Dis* 2003; 5: 1-7.
- 58) GÁMEZ J, PLAYÁN A, ANDREU AL, BRUNO C, NAVARRO C, CERVERA C, ARBÓS MA, SCHWARTZ S, ENRIQUEZ JA, MONTOYA J. Familial multiple symmetric lipomatosis associated with the A8344G mutation of mitochondrial DNA. *Neurology* 1998; 51: 258-260.
- 59) MANCUSO M, ORSUCCI D, ANGELINI C, BERTINI E, CARELLI V, COMI GP, MINETTI C, MOGGIO M, MONGINI T, SERVIDEI S, TONIN P, TOSCANO A, UZIEL G, BRUNO C, CALDARAZZO IENCO E, FILOSTO M, LAMPERTI C, MARTINELLI D, MORONI I, MUSUMECI O, PEGORARO E, RONCHI D, SANTORELLI FM, SAUCHELLI D, SCARPELLI M, SCIACCO M, SPINAZZI M, VALENTINO ML, VERCELLI L, ZEVIANI M, SICILIANO G. Phenotypic heterogeneity of the 8344A>G mtDNA "MERRF" mutation. *Neurology* 2013; 80: 2049-2054.
- 60) PERERA U, KENNEDY BA, HEGELE RA. Multiple symmetric lipomatosis (Madelung disease) in a large Canadian family with the mitochondrial MTTK c.8344A>G variant. *J Investig Med High Impact Case Rep* 2018; 6: 2324709618802867.
- 61) CAPEL E, VATIER C, CERVERA P, STOJKOVIC T, DISSE E, COTTEREAU AS, AUCLAIR M, VERPONT MC, MOSBAH H, GOURDY P, BARRAUD S, MIOUET A, ZÜCHNER S, BONNEFOND A, FROGUEL P, CHRISTIN-MAITRE S, DELEMER B, FÈVE B, LAVILLE M, ROBERT J, TENENBAUM F, LASCOLS O, VIGOUROUX C, JÉRU I. MFN2-associated lipomatosis: clinical spectrum and impact on adipose tissue. *J Clin Lipidol* 2018; 12: 1420-1435.
- 62) HERBST KL, TANNOCK LR, DEEB SS, PURNELL JQ, BRUNZELL JD, CHAIT A. Köbberling type of familial partial lipodystrophy: an underrecognized syndrome. *Diabetes Care* 2003; 26: 1819-1824.
- 63) LIGHTBOURNE M, BROWN RJ. Genetics of lipodystrophy. *Endocrinol Metab Clin North Am* 2017; 46: 539-554.
- 64) BROEKEMA MF, SAVAGE DB, MONAJEMI H, KALKHOVEN E. Gene-gene and gene-environment interactions in lipodystrophy: lessons learned from natural PPARG mutants. *Biochim Biophys Acta Mol Cell Biol Lipids* 2019; 1864: 715-732.
- 65) SHEARIN AL, MONKS BR, SEALE P, BIRNBAUM MJ. Lack of AKT in adipocytes causes severe lipodystrophy. *Mol Metab* 2016; 5: 472-479.
- 66) GARG A, SANKELLA S, XING C, AGARWAL AK. Whole-exome sequencing identifies ADRA2A mutation in atypical familial partial lipodystrophy. *JCI Insight* 2016; 1(9). pii: e86870.
- 67) GESTA S, BLÜHER M, YAMAMOTO Y, NORRIS AW, BERNDT J, KRALISCH S, BOUCHER J, LEWIS C, KAHN CR. Evidence for a role of developmental genes in the origin of obesity and body fat distribution. *Proc Natl Acad Sci U S A* 2006; 103: 6676-6681.
- 68) HEID IM, JACKSON AU, RANDALL JC, WINKLER TW, QI L, STEINTHORSDDOTTIR V, THORLEIFSSON G, ZILLIKENS MC, SPELIOTES EK, MÄGI R, WORKALEMAHU T, WHITE CC, BOUATIA-NAJI N, HARRIS TB, BERNDT SI, INGELSSON E, WILLER CJ, WEEDON MN, LUAN J, VEDANTAM S, ESKO T, KILPELÄINEN TO, KUTALIK Z, LI S, MONDA KL, DIXON AL, HOLMES CC, KAPLAN LM, LIANG L, MIN JL, MOFFATT MF, MOLONY C, NICHOLSON G, SCHATZ EE, ZONDERVAN KT, FEITOSA MF, FERREIRA T, LANGO ALLEN H, WEYANT RJ, WHEELER E, WOOD AR, MAGIC, ESTRADA K, GODDARD ME, LETTRE G, MANGINO M, NYHOLT DR, PURCELL S, SMITH AV, VISSCHER PM, YANG J, MCCARROLL SA, NEMESH J, VOIGHT BF, ABSHER D, AMIN N, ASPELUND

- T, COIN L, GLAZER NL, HAYWARD C, HEARD-COSTA NL, HOTTENGA JJ, JOHANSSON A, JOHNSON T, KAAKINEN M, KAPUR K, KETKAR S, KNOWLES JW, KRAFT P, KRAJA AT, LAMINA C, LEITZMANN MF, MCKNIGHT B, MORRIS AP, ONG KK, PERRY JR, PETERS MJ, POLASEK O, PROKOPENKO I, RAYNER NW, RIPATTI S, RIVADENEIRA F, ROBERTSON NR, SANNA S, SOVIO U, SURAKKA I, TEUMER A, VAN WING-ERDEN S, VITART V, ZHAO JH, CAVALCANTI-PROENÇA C, CHINES PS, FISHER E, KULZER JR, LECOEUR C, NARISU N, SANDHOLT C, SCOTT LJ, SILANDER K, STARK K, TAMMESOO ML, TESLOVICH TM, TIMPSON NJ, WATANABE RM, WELCH R, CHASMAN DI, COOPER MN, JANSSON JO, KETTUNEN J, LAWRENCE RW, PELLIKKA N, PEROLA M, VANDENPUT L, ALAVERE H, ALMGREN P, ATWOOD LD, BENNETT AJ, BIFFAR R, BONNYCASTLE LL, BORNSTEIN SR, BUCHANAN TA, CAMPBELL H, DAY IN, DEI M, DÖRR M, ELLIOTT P, ERDOS MR, ERIKSSON JG, FREIMER NB, FU M, GAGET S, GEUS EJ, GJESING AP, GRALLERT H, GRÄSSLER J, GROVES CJ, GUIDUCCI C, HARTIKAINEN AL, HASSANALI N, HAVULINNA AS, HERZIG KH, HICKS AA, HUI J, IGL W, JOUSILAHTI P, JULA A, KAJANTIE E, KINNUNEN L, KOLCIC I, KOSKINEN S, KOVACS P, KROEMER HK, KRZEL V, KUUSISTO J, KVALOY K, LAITINEN J, LANTIERI O, LATHROP GM, LOKKI ML, LUBEN RN, LUDWIG B, MCARDLE WL, MCCARTHY A, MORKEN MA, NELIS M, NEVILLE MJ, PARÉ G, PARKER AN, PEDEN JF, PICHLER I, PIETILÄINEN KH, PLATOU CG, POUTA A, RIDDERSTRALE M, SAMANI NJ, SARAMEES J, SINISALO J, SMIT JH, STRAWBRIDGE RJ, STRINGHAM HM, SWIFT AJ, TEDER-LAVING M, THOMSON B, USALA G, VAN MEURS JB, VAN OMMEN GJ, VATIN V, VOLPATO CB, WALLASCHOFSKI H, WALTERS GB, WIDEN E, WILD SH, WILLEMSEN G, WITTE DR, ZGAGA L, ZITTING P, BEILBY JP, JAMES AL, KÄHÖNEN M, LEHTIMÄKI T, NIEMINEN MS, OHLSSON C, PALMER LJ, RAITAKARI O, RIDKER PM, STUMVOLL M, TÖNJES A, VIKARI J, BALKAU B, BEN-SHLOMO Y, BERGMAN RN, BOEING H, SMITH GD, EBRAHIM S, FROGUEL P, HANSEN T, HENGSTENBERG C, HVEEM K, ISOMAA B, JØRGENSEN T, KARPE F, KHAW KT, LAAKSO M, LAWLOR DA, MARRE M, MEITINGER T, METSPALU A, MIDTHJELL K, PEDERSEN O, SALOMAA V, SCHWARZ PE, TUOMI T, TUOMILEHTO J, VALLE TT, WAREHAM NJ, ARNOLD AM, BECKMANN JS, BERGMANN S, BOERWINKLE E, BOOMSMA DI, CAULFIELD MJ, COLLINS FS, EIRIKSDOTTIR G, GUDNASON V, GYLLENSTEN U, HAMSTEN A, HATTERSLEY AT, HOFMAN A, HU FB, ILLIG T, IRIBARREN C, JARVELIN MR, KAO WH, KAPRIO J, LAUNER LJ, MUNROE PB, OOSTRA B, PENNINX BW, PRAMSTALLER PP, PSATY BM, QUERTERMOS T, RISSANEN A, RUDAN I, SHULDINER AR, SORANZO N, SPECTOR TD, SYVANEN AC, UDA M, UITTERLINDEN A, VÖLZKE H, VOLLENWEIDER P, WILSON JF, WITTEMAN JC, WRIGHT AF, ABECASIS GR, BOEHNKE M, BORECKI IB, DELOUKAS P, FRAYLING TM, GROOP LC, HARITUNIANI T, HUNTER DJ, KAPLAN RC, NORTH KE, O'CONNELL JR, PELTONEN L, SCHLESSINGER D, STRACHAN DP, HIRSCHHORN JN, ASSIMES TL, WICHMANN HE, THORSTEINSDOTTIR U, VAN DUJIN CM, STEFANSSON K, CUPPLES LA, LOOS RJ, BARROSO I, MCCARTHY MI, FOX CS, MOHLKE KL, LINDGREN CM. Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nat Genet* 2010; 42: 949-960.
- 69) CHU AY, DENG X, FISHER VA, DRONG A, ZHANG Y, FEITOSA MF. Multiethnic genome-wide meta-analysis of ectopic fat depots identifies loci associated with adipocyte development and differentiation. *Nat Genet* 2017; 49: 125-130.
- 70) MÄKINEN T, JUSSILA L, VEIKKOLA T, KARPANEN T, KETTUNEN MI, PULKKANEN KJ, KAUPPINEN R, JACKSON DG, KUBO H, NISHIKAWA S, YLÄ-HERTTUALA S, ALITALO K. Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice expressing soluble VEGF receptor-3. *Nat Med* 2001; 7: 199-205.
- 71) MENDOLA A, SCHLÖGEL MJ, GHALAMKARPOUR A, IRRTHUM A, NGUYEN HL, FASTER E, BYGUM A, VAN DER VLEUTEN C, FAGERBERG C, BASELGA E, QUERE I, MULLIKEN JB, BOON LM, BROUILLARD P, VIKKULA M; Lymphedema Research Group. Mutations in the VEGFR3 signaling pathway explain 36% of familial lymphedema. *Mol Syndromol* 2013; 4: 257-266.
- 72) COHEN P, LEVY JD, ZHANG Y, FRONTINI A, KOLODIN DP, SVENSSON KJ, LO JC, ZENG X, YE L, KHANDEKAR MJ, WU J, GUNAWARDANA SC, BANKS AS, CAMPOREZ JP, JURCZAK MJ, KAJIMURA S, PISTON DW, MATHIS D, CINTI S, SHULMAN GI, SEALE P, SPIEGELMAN BM. Ablation of PRDM16 and beige adipose causes metabolic dysfunction and a subcutaneous to visceral fat switch. *Cell* 2014; 156: 304-316.
- 73) KUANG J, ZHANG Y, LIU Q, SHEN J, PU S, CHENG S, CHEN L, LI H, WU T, LI R, LI Y, ZOU M, ZHANG Z, JIANG W, XU G, QU A, XIE W, HE J. Fat-specific Sirt6 ablation sensitizes mice to high-fat diet-induced obesity and insulin Resistance by inhibiting lipolysis. *Diabetes* 2017; 66: 1159-1171.
- 74) MARTÍNEZ-JIMÉNEZ V, CORTEZ-ESPINOZA N, RODRÍGUEZ-VARELA E, VEGA-CÁRDENAS M, BRIONES-ESPINOZA M, RUIZ-RODRÍGUEZ VM, LÓPEZ-LÓPEZ N, BRISEÑO-MEDINA A, TURJÁN-ESPINOZA E, PORTALES-PÉREZ DP. Altered levels of sirtuin genes (SIRT1, SIRT2, SIRT3 and SIRT6) and their target genes in adipose tissue from individual with obesity. *Diabetes Metab Syndr* 2019; 13: 582-589.
- 75) LI H, WU G, FANG Q, ZHANG M, HUI X, SHENG B, WU L, BAO Y, LI P, XU A, JIA W. Fibroblast growth factor 21 increases insulin sensitivity through specific expansion of subcutaneous fat. *Nat Commun* 2018; 9: 272.
- 76) PAZOS P, LIMA L, TOVAR S, GONZÁLEZ-TOUCEDA D, DIÉGUEZ C, GARCÍA MC. Divergent responses to thermogenic stimuli in BAT and subcutaneous adipose tissue from interleukin 18 and interleukin 18 receptor 1-deficient mice. *Sci Rep* 2015; 5: 17977.
- 77) LEE VR, BARR KJ, KELLY JJ, JOHNSTON D, BROWN CFC, ROBB KP, SAYEDYAHOSSEIN S, HUANG K, GROS R, FLYNN LE, PENUELA S. Pannexin 1 regulates adipose stromal cell differentiation and fat accumulation. *Sci Rep* 2018; 8: 16166.
- 78) MIKA A, KASKA L, KORCZYNSKA J, MIROWSKA A, STEPNOWSKI P, PROCZKO M, RATNICKI-ŚKLUCKI K, GOYKE E, SLEDZINSKI T. Visceral and subcutaneous adipose tissue stearoyl-CoA desaturase-1 mRNA levels and fatty acid desaturation index positively correlate with BMI in morbidly obese women. *Eur J Lipid Sci Technol* 2015; 117: 926-932.
- 79) YEE JK, PHILLIPS SA, ALLAMEHZADEH K, HERBST KL. Subcutaneous adipose tissue fatty acid desaturation in adults with and without rare adipose disorders. *Lipids Health Dis* 2012; 11: 19.
- 80) HARVEY NL, SRINIVASAN RS, DILLARD ME, JOHNSON NC, WITTE MH, BOYD K, SLEEMAN MW, OLIVER G. Lymphatic vascular defects promoted by Prox1 haploinsufficiency cause adult-onset obesity. *Nat Genet* 2005; 37: 1072-1081.

- 81) SHAO M, HEPLER C, VISHVANATH L, MACPHERSON KA, BUSBUSO NC, GUPTA RK. Fetal development of subcutaneous white adipose tissue is dependent on Zfp423. *Mol Metab* 2016; 6: 111-124.
- 82) AMANN-VESTI BR, FRANZECK UK, BOLLINGER A. Microlymphatic aneurysms in patients with lipedema. *Lymphology* 2001; 34: 170-175.
- 83) TODD M. Lipoedema: presentation and management. *Br J Community Nurs* 2010; 15: S10-S16.
- 84) CRESCENZI R, MARTON A, DONAHUE PMC, MAHANY HB, LANTS SK, WANG P, BECKMAN JA, DONAHUE MJ, TITZE J. Tissue sodium content is elevated in the skin and subcutaneous adipose tissue in women with lipedema. *Obesity (Silver Spring)* 2018; 26: 310-317.
- 85) WIIG H, LUFT FC, TITZE JM. The interstitium conducts extrarenal storage of sodium and represents a third compartment essential for extracellular volume and blood pressure homeostasis. *Acta Physiol (Oxf)* 2018; 222.
- 86) MICHELINI S, FIORENTINO A, CESTARI M, FANTEGROSSI MR, CARDONE M. Lipedema and metabolic disease. *Oedeem Oncologie* 2018; 3: 38-39.
- 87) SACEY E, OSTERGAARD P. Genetic research on lipedema at St George's University of London, UK. *Oedeem Oncologie* 2018; 3: 9-10.