

This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

Hit-to-lead optimization of mouse Trace Amine Associated Receptor 1 (mTAAR1) agonists with a diphenylmethane-scaffold: Design, Synthesis, and biological study

Journal:	<i>Journal of Medicinal Chemistry</i>
Manuscript ID	jm-2016-01092q.R1
Manuscript Type:	Article
Date Submitted by the Author:	07-Sep-2016
Complete List of Authors:	Chiellini, Grazia; Università di Pisa, Department of Pathology Nesi, Giulia; Università di Pisa, Dipartimento di Farmacia Sestito, Simona; Università di Pisa, Dipartimento di Farmacia Chiarugi, Sara; Università di Pisa, Dipartimento di Farmacia Runfola, Massimiliano; Università di Pisa, Dipartimento di Farmacia Espinoza, Stefano; Istituto Italiano di Tecnologia, Dept of Neuroscience and Brain Technologies Sabatini, Martina; Università di Pisa, Department of Pathology Bellusci, Lorenza; Università di Pisa, Department of Pathology Laurino, Annunziata; università degli studi di firenze, neurofarba Cichero, Elena; University of genoa, Pharmacy Gainetdinov, Raul; Istituto Italiano di Tecnologia, NBT; Institute of Translational Biomedicine, , St. Petersburg State University; Skolkovo Institute of Science and Technology Fossa, Paola; University of Genova, Farmacia Raimondi, Laura; università degli studi di firenze, neurofarba Zucchi, Riccardo; Università di Pisa, Dept of Pathology Rapposelli, Simona; Università di Pisa, Dipartimento di Farmacia

SCHOLARONE™
Manuscripts

1
2
3 **Hit-to-lead optimization of mouse Trace Amine Associated Receptor 1 (mTAAR1) agonists**
4
5 **with a diphenylmethane-scaffold: Design, Synthesis, and biological study**
6
7
8
9

10
11 Grazia Chiellini^{1*}, Giulia Nesi², Simona Sestito², Sara Chiarugi², Massimiliano Runfola², Stefano
12 Espinoza³, Martina Sabatini¹, Lorenza Bellusci¹, Annunziata Laurino⁴, Elena Cichero⁵, Raul. R.
13 Gainetdinov^{6,7}, Paola Fossa⁵, Laura Raimondi⁴, Riccardo Zucchi¹, Simona Rapposelli^{2*}.
14
15
16

17
18 *¹Dept. of Pathology, University of Pisa, Pisa, Italy, ²Dept. of Pharmacy, University of Pisa, Pisa,*
19 *Italy, ³Department of Neuroscience and Brain Technologies, Istituto Italiano di Tecnologia,*
20 *Genova, Italy, ⁴Dept. of NEUROFARBA; Section of Pharmacology, University of Florence,*
21 *Italy, ⁵Dept. of Pharmacy, University of Genoa, Genoa, Italy. ⁶Institute of Translational*
22 *Biomedicine, St. Petersburg State University, St. Petersburg, 199034, Russia, ⁷Skolkovo Institute of*
23 *Science and Technology (Skoltech) Skolkovo, Moscow region, 143025, Russia.*
24
25
26
27
28
29
30
31
32
33
34
35

36 **Keywords:** mTAAR1 agonist; diphenylmethane-scaffold; Thyronamine; T1AM; T0AM
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract.

The trace amine-associated receptor 1 (TAAR1) is a G-protein-coupled receptors (GPCR) potentially activated by a variety of molecules besides trace amines (TAs), including thyroid hormone-derivatives like 3-iodothyronamine (T1AM), catechol-O-methyltransferase products like 3-methoxytyramine, and amphetamine-related compounds. Accordingly, TAAR1 is considered a promising target for medicinal development. To gain more insights into TAAR1 physiological functions and validation of its therapeutic potential we recently developed a new class of thyronamine-like derivatives. Among them compound SG2 showed high affinity and potent agonist activity at mouse TAAR1. In the present work we describe design, the synthesis and SAR study of a new series of compounds (**1-16**) obtained by introducing specific structural changes at key points of our lead-compound SG2 skeleton. Five of the newly synthesized compounds displayed mTAAR1 agonist activity higher than both SG2 and T1AM. Selected diphenylmethane analogs, namely **1** and **2**, showed potent functional activity in *in vitro* and *in vivo* models.

Introduction.

Thyronamine (TAM) is generally used to indicate a class of endogenous compounds deriving from thyroid hormones through metabolic reactions such as deiodination and decarboxylation¹. Over the past decade, thyronamines have been recognized as ligands of trace amine - associated receptor 1 (TAAR1), and to date they represent a thriving research field. Currently, only two compounds that belong to the class of TAM have been identified *in vivo*: the 3-iodothyronamine (T1AM) and the thyronamine (T0AM). These compounds have been postulated to derive from thyroid hormone (T4) through deiodination and decarboxylation. T1AM is widely expressed in various tissues, such as brain, liver, heart and blood and it seems to be the most abundant thyronamine present in circulation².

Among several potential receptors, it has been proved that T1AM binds to TAAR1, a G-protein coupled receptor (GPCR) that has been identified in specific areas of the central nervous system and in some peripheral areas. The binding of T1AM to TAAR1 receptor leads to increase of cAMP levels by the activation of the adenylate cyclase. Further studies also highlighted that T1AM interact with other targets such as the biogenic amines transporters and the apoB100 protein. Thyronamines, and in particular T1AM, produce different functional effects, although their physiological role still remains unclear. Among the most significant effects attributed to T1AM are a hypothermic as well as negative inotropic and chronotropic cardiac effects. Moreover, endocrine and metabolic effects such as the modulation of insulin secretion, the inhibition of catecholamines resorption at the neuronal level and an increased metabolism of lipids at the expense of that of carbohydrates have been also described³. The wide variety of functions attributed to T1AM makes this molecule a potentially useful tool for the treatment of many diseases such as obesity, neuropsychiatric disorders and cancer. Unfortunately, TAMs are rapidly metabolized by different enzyme systems, such as amino-oxidase (MAO, SSAO), deiodinase (DIO3), sulfotransferase (SULT1A1 and SULT1A3), N-acetyltransferase, and glucuronidase. The action of these enzymes is

1
2
3 a limit to their therapeutic use. Consequently, the great therapeutic potential as well as the lack of
4
5 new useful tools to elucidate the physiological function of T1AM, have pushed medicinal chemists
6
7 to develop new synthetic analogues of T1AM.
8
9

10 Recently, with the aim of increasing the number of selective ligands for TAAR1 receptor and
11
12 provide new tools to facilitate the understanding of the physiological functions of this receptor, we
13
14 reported the design and synthesis of a new class of thyronamine-like compounds with a
15
16 diphenylmethane scaffold, which proved to be synthetically more accessible than endogenous
17
18 T1AM and T0AM^{4, 5}. Within the small series of analogues previously synthesized, SG1 and SG2
19
20 were found to activate mouse TAAR1 receptor (mTAAR1) with a potency comparable to the
21
22 corresponding endogenous ligands⁴ (Figure 1). In particular, SG2, the analogue of T1AM, showed
23
24 an $EC_{50} = 240$ nM comparable to T1AM ($EC_{50} = 189$ nM). In order to expand the SAR study for
25
26 this new class of compounds, in the present work we describe the design and synthesis of
27
28 compounds **1-16** obtained by introducing specific structural changes at key points of our lead-
29
30 compound SG2 skeleton. The structural optimization process of the new thyronamine-like
31
32 compounds was based on molecular docking studies that highlighted the main differences in the
33
34 interaction between the endogenous ligand T1AM or the synthetic analogue SG2 with the mTAAR1
35
36 receptor binding site. The newly designed compounds were first screened *in vitro* for mTAAR1
37
38 activation. Five compounds (**1,2,4,7** and **8**) showed a higher potency than the lead molecule SG2, as
39
40 well as, the endogenous ligand T1AM, with **1** and **2** being the most potent of the new series. These
41
42 compounds were then evaluated both *in vitro* and *in vivo* to characterize their ability to modulate
43
44 plasma glucose levels.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Structure of T1AM and the lead-molecules SG1 and SG2 with a diphenylmethane molecular scaffold

Results.

Design of new thyronamine-like analogues.


In our previous work, we explored the pharmacological profile of a new series of T₁AM diphenylmethane analogues, and among these compounds SG1 and SG2 were found to be a good mimic of the corresponding endogenous ligands T0AM and T1AM⁴, respectively. In addition, a docking study was performed to investigate the main differences between T1AM and the lead diphenylmethane-analogue SG2 on mTAAR1 receptor binding. According to this study, T1AM appears to be highly stabilized in the binding site through the formation of two H-bonds between the protonated amino group of the ethylamine side chain and aspartic acid 102 (D102) and tyrosine 291 (Y291). Furthermore, T1AM established an additional H-bond between the hydroxyl-group on the outer ring and arginine 82 (R82), while no significant amino acid interactions were observed for the biaryl-etheral oxygen of T1AM. Notably, the SG2 derivative appeared to share most of the main mTAAR1/T1AM interactions⁴. Interestingly, the replacement of the hydroxyl group on the T1AM outer ring with an amino group, as in SG2, allowed the formation of a cation- π interaction, while no H-bonds to the R82 backbone oxygen atom were observed. Conceivably, the oxygen atom in the SG2-inner ring side chain may induce a shift of the molecule within the binding site, thus preventing the H-bond formation with R82. In order to expand the SAR study for this new class of

1
2
3 thyronamine analogues, we performed specific structural changes at key points of the SG2-
4
5 molecular scaffold. Initially, to restore the H-bond with R82, the oxo-ethylamino side chain of SG2
6
7 was replaced with an ethylamine one (**1-6**). Among these new compounds, analogue **2** was
8
9 synthesized to evaluate the effects induced by the recovery of the phenolic function as in T1AM.
10
11 Additionally, to gain further insights on how the pattern of substitutions modulate the SG-analogues
12
13 potency as mTAAR1 agonists, we investigated the following structure modifications: (a) the
14
15 introduction of small alkyl substituents (Me, i-Pr) on both outer- and inner-rings (**4,6-10**); (b) the
16
17 removal of the methylene-bridge, producing the diaryl-analogues **5** and **6**; (c) the introduction of a
18
19 biguanide-moiety at the oxy-alkyl side chain (**15,16**) and finally (d) mono- and bis-N-acetylation (**3**,
20
21 **9-15**). This last chemical modification has been suggested by a recent study conducted by Hoefig et
22
23 al⁶, showing that O-acetyl and N-acetyl derivatives of T1AM may be found in white adipose tissue
24
25 and liver endogenous cells. Notably, N-Ac-T1AM was also found to be an endogenous cardioactive
26
27 metabolite⁶. On the basis of these findings, the corresponding mono- and bis-N-acetylated
28
29 analogues of SG1 and SG2 lead compounds (i.e. **11,12** and **13,14**, respectively) as well as the new
30
31 acetylated products namely **3**, **9,10** and **15**, were synthesised.
32
33
34
35
36

37 **Synthesis.**

38
39
40 The derivatives **1-6** in which the oxo-ethylamino side chain of SG2 has been replaced with the
41
42 ethylamine one have been synthesized as reported in Scheme 1. Briefly, the products **17-20**
43
44 obtained by the palladium(0)-catalyzed Suzuki–Miyaura cross-coupling reaction of the
45
46 trifluoroborate salt or boronic acid with the appropriate benzylbromide⁴, reacted with SOCl₂ to give
47
48 benzylchloride **21-24**. Subsequently, the nucleophilic substitution with NaCN afforded the
49
50 compound **25-28** with high yields. As previously described⁴, reduction of nitrile derivative with
51
52 LiAlH₄ in the presence of a Lewis's acid afforded the diamine derivative **1,4-6**. Then, the treatment
53
54 of **1** with a mixture of Ac₂O/NaHCO₃ provided the acetylanilide **3** (Scheme 1a).
55
56
57
58
59
60

Scheme 1




Reagents and conditions: **a:** SOCl_2 , CHCl_3 , rt, 2h; **b:** NaCN , $\text{H}_2\text{O}/\text{CH}_3\text{CN}$, mw; **c:** LiAlH_4 , AlCl_3 , THF, reflux, 12h; **d:** $\text{Ac}_2\text{O}/\text{NaHCO}_3$, rt, 1h, **e:** H_2 , Pd-C, AcOH, 12h; **f:** NaNO_2 , H_2SO_4 , H_2O , 100°C , 1h.

Compound **2** was obtained starting from the acetonitrile derivative **25**. Reduction of the nitro-group with H_2 in the presence of 10% Pd-C afforded the aniline **29**. The subsequent hydrolysis of the diazonium salt obtained by reaction of **29** with $\text{NaNO}_2/\text{H}_2\text{SO}_4$ afforded the final product **2** (Scheme 1b).

The Suzuki Miyaura cross-coupling reaction of the 2,6-dimethyl-4-methoxy-boronic acid with the 4-nitrobenzyl bromide gave **31** which was demethylated with BBr_3 at 0°C affording to phenol **32**. The subsequent reaction of α -bromide-acetonitrile with **32** followed by reduction to the corresponding diamine-derivative gave the superior homolog of SG2 (**7**) (Scheme 2).

Scheme 2




Reagents and conditions: a: 4-Nitrobenzyl bromide, K_2CO_3 , PdCl_2 , Acetone/ H_2O , rt, 72 h; b: BBr_3 , DCM, 0°C , 1 h; c: BrCH_2CN , DMF, Cs_2CO_3 , rt, 30'; d: LiAlH_4 , AlCl_3 , THF, reflux, 12 h.

Derivative **8-10** substituted with an *i*-Pr group in the outer ring and with a small *N*-alkyl (**8**) or *N*-acetyl-group (**9,10**) were synthesized starting from the cross coupling reaction of *N*-(4-(bromomethyl)-2-isopropylphenyl)acetamide (**34**) (see SI, Scheme S1) and the (4-(hydroxymethyl)-2-methylphenyl)fluoroboronate salt (Scheme 3). The tandem reactions with SOCl_2 and NaCN gave the nitrile derivative **37** with high yields. Subsequent reduction with $\text{LiAlH}_4/\text{AlCl}_3$ afforded the product **8**, which was then reacted with Ac_2O in the presence of NaHCO_3 in a ratio of 1:1 or 1:2 to give **9** and **10**, respectively.

As concerns the synthesis of the mono- and *N*-diacetylated derivatives **11-14** and the biguanide analogue of SG2 (**15** and **16**), the synthetic pathway as well as the procedures are reported in SI (Scheme S2).

Moreover, the addition of small alkyl groups in the inner ring of SG2 (**7**), as well as in the outer ring of the N-ethyl analogue of **1** (**8**), turned out to be successful, thus obtaining new compounds with a potency 2-fold higher than the lead compound SG2.


Table 1. Activity of the new diphenylmethane derivatives **1-6, 8-10** using BRET-based assay in mTAAR1 transfected HEK 293 cells. The EC₅₀ values are expressed in nM.





1-6;8-10

compd	X	n	R	R ₁	R ₂	Y	mTAAR1 EC ₅₀ (nM)
1	NH ₂	1	Me	H	H	NH ₂	119
2	OH	1	Me	H	H	NH ₂	98
3	NHAc	1	Me	H	H	NH ₂	3100
4	NH ₂	1	Me	Me	H	NH ₂	158
5	NH ₂	0	Me	H	H	NH ₂	1180
6	NH ₂	0	Me	Me	H	NH ₂	633
8	NHEt	1	Me	H	i-Pr	NH ₂	102
9	AcNEt	1	Me	H	i-Pr	NH ₂	440
10	AcNEt	1	Me	H	i-Pr	NHAc	>10000

Table 2. Activity of the new diphenylmethane derivatives **7**, **11-16** using BRET-based assay in mTAAR1 transfected HEK 293 cells. The EC₅₀ values are expressed in nM.



7,11-16

compd	X	n	R	R ₁	Y	mTAAR1 EC ₅₀ (nM)
7	NH ₂	1	Me	Me	NH ₂	134
11	NHAc	1	H	H	NH ₂	>10000
12	NHAc	1	H	H	NHAc	>10000
13	NHAc	1	Me	H	NH ₂	>10000
14	NHAc	1	Me	H	NHAc	>10000
15	NHAc	1	Me	H		>10000
16	NH ₂	1	Me	H		>10000




Figure 2. Dose-response curves of the most active SG2-analogues. All the compounds are full agonists (Efficacy = 100%)

Docking study.

In our previous work, we investigated T₁AM and a series of T₁AM analogues whose potency profile proved to be highly related to the presence of a proper basic feature linked to an aromatic core⁴. Indeed, our molecular modelling studies pointed out the relevance of a key salt bridge between the protonated amine group of SG1 and SG2 lead compounds and the highly conserved residue D102, together with a number of π - π stacking and cation- π contacts with the surrounding amino acids, as we previously detected in other series of TAAR1 ligands^{8,9}.

These findings were supported by further homology modelling and docking studies we performed within the murine and human TAAR1 and TAAR5 receptors with respect to T₁AM ligand¹⁰, and were also in accordance with the key role played by a proper basic feature included in any TAAR1

1
2
3 ligand, as reported in literature¹¹. Notably, this information was definitively validated by the
4
5 mutagenesis data involving the D102 residue, reported by Reese¹².
6

7 In order to rationalize the pharmacological results and refine the computational models, *in silico*
8
9 docking studies on the most active compounds (**1,2,7** and **8**) were also performed, analysing their
10
11 interactions with the putative receptor binding site (see SI, Table S1).
12

13 In agreement with the experimental data, our docking calculations revealed that the presence of the
14
15 ethylamine chain appears to be the most effective choice, in particular within those series of
16
17 compounds bearing a mono methyl-substituted inner ring (**1,2** and **8**). Based on a comparison with
18
19 the prototype T1AM, the most promising newly designed analogues were overturned within the
20
21 receptor binding site, maintaining in any case the key contact with D102 which is required for
22
23 TAAR1 agonism (see SI, Figure S1)¹².
24
25
26

27 Notably, the most effective compounds **1** and **2** were characterized by a reversed docking mode
28
29 when compared to that previously described for SG2⁴. Briefly, we originally highlighted the key
30
31 role played by the protonated amino group of the ethoxyamine side chain of SG2 in H-bonding
32
33 with D102 and Y291, while the inner-phenyl group was engaged in π - π stacking with Y287. On the
34
35 contrary, **1** (and **2**) displayed one H-bond between the aniline portion in the outer ring (or the
36
37 hydroxyl group) and the D102 negative-charged side-chain, while the protonated basic side chain
38
39 appeared to be involved in one H-bond and in a salt-bridge with T83 and D284, respectively, as
40
41 shown in Figure 3a and 3b, respectively.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3. Docking mode of compounds SG2 (C atom cyan) and (a) **1** (C atom deep magenta) (b) **2** (C atom; coral) (c) **7** (C atom; light green) and (d) **8** (C atom; yellow) at the murine TAAR1 binding site. The most important residues are labelled.

The addition of a methyl-group in the inner-ring of SG2 led to a 2-fold increase of potency for **7**, with an EC_{50} of 134nM vs 240 nM for SG2. The docking study showed that also **7** takes an overturned positioning with respect to SG2 in the binding site of mTAAR1 (Figure 3c).

1
2
3 Presumably, the presence of an additional substituent into the inner ring causes a shift of the
4 molecule within the binding site promoting the interaction of the aniline nitrogen atom in the outer
5 ring with D102 residue, while the side chain's nitrogen atom interacts with D284.
6
7


8
9 Finally, the introduction of small alkyl groups into the outer ring of **1** as in compound **8** didn't
10 prevent the interaction between the nitrogen atom of aniline with D102 residue. Indeed, the EC₅₀
11 value of **8** (EC₅₀ = 102nM) was comparable to that of **1** (EC₅₀ = 119nM).
12
13
14
15
16
17
18

19 **Modulation of hepatic glucose production.**

20
21
22 The hyperglycemic effect observed after administration of exogenous T1AM had previously been
23 attributed only to a modulation of insulin and/or glucagone secretion¹³⁻¹⁵. Recently, our study¹⁶
24 showed that in human hepatocarcinoma cells (HepG2) T1AM treatment directly stimulated glucose
25 production (EC₅₀ = 0.84 μM), if adequate gluconeogenic substrates were provided. Interestingly,
26 the effect of T1AM on gluconeogenesis displayed a bell-shaped dose-response curve, suggesting a
27 biphasic effect of T1AM on the modulation of gluconeogenesis. In the present study, the effects
28 induced by two of the most potent new mTAAR1 agonists (i.e. **1** and **2**) on gluconeogenesis were
29 assessed. Preliminarily, HepG2 cells were incubated with each compound at three different doses
30 (0.1, 1 and 10μM). By comparison, similar experiments were performed using T1AM. As shown in
31 Figure 4a, incubation with 0.1 and 1 μM of compound **2** induced a significant increase in glucose
32 production (P<0.01), while a dose of 10μM induced a decreased stimulation of glucose production.
33
34 On the other hand, compound **1** showed a dose-dependent increase of glucose production, which
35 reached statistical significance only when compound **1** was used at the highest dose (10μM). As
36 shown in Figure 4a, treatment of HepG2 cells with T1AM (0.1, 1 and 10 μM) confirmed the bell-
37 shaped dose-response curve observed in our previous study¹⁶. To better define the dose dependence
38 of **1** and **2** effects on glucose production additional experiments were carried out using a wider
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

range of doses for both compounds. As shown in Figure 4b the EC₅₀ calculated for the effect on gluconeogenesis was in the submicromolar range for **2**, and in the supramicromolar range for **1**.

Figure 4: (a) Glucose production in HepG2 cell cultures that were incubated for 4 h in glucose production buffer after adding vehicle (DMSO) or compounds **1**, **2** and **T1AM**. Results are expressed as mean \pm S.E.M. and are normalized to the total cell protein content determined in cell lysates. (n=3 in each case). **(b)** Glucose production in HepG2 cell in response to increasing concentrations of compound **1** (0.1, 1, 10 and 20 μ M) and **2** (0.001, 0.01, 0.1, 1 and 10 μ M). Results are mean \pm S.E.M. and are normalized to the total cell protein content determined in cell lysates. (n=6 in each case). *P<0.05, **P<0.01, ***P<0.001 by ANOVA.




Modulation of plasma glucose level.

As previously reported by us thyronamine-like analogues SG1 and SG2 demonstrated to be able to increase plasma glycaemia with a potency comparable to that of the corresponding endogenous thyronamine (i.e. T0AM and T1AM)⁴. Thus, in the present work the effects induced by **1** and **2** derivatives on plasma glucose level were also assessed.

Figure 5 shows that a single low dose (4.0 μ g/kg, i.p.) of **1** or **2** significantly increases plasma glycaemia in CD-1 mice with a potency comparable to that previously shown by lead-compound SG2⁴.

Figure 5. Effect of **1** and **2** on plasma glycaemia ($\text{mg}\cdot\text{L}^{-1}$) in CD1-mice. Plasma glucose was measured 15 min after test compound or vehicle injection (i.p.) in blood collected from the tail veins of 4 h starved mice ($n = 8$ in each group). Results are expressed as means $\text{mean}\pm\text{SEM}$. * $P < 0.05$ vs vehicle. *** $P < 0.001$ vs vehicle.



Conclusion

Our ongoing efforts to develop thyronamine-like analogues synthetically more accessible than endogenous thyronamines yielded a new class of potent mTAAR1 diphenylmethane ligands.

A medicinal chemistry optimization combined to docking studies were used starting from the molecular scaffold of our recently developed SG-compounds to design the new ligands. In particular, we investigated: (i) the replacement of the oxyethylamino side chain of SG2 with an ethylamino one (ii) the introduction of small alkyl substituents (Me, i-Pr) on both outer- and inner-rings and (iii) the recovery of the phenolic function as in T1AM. Overall these chemical modifications resulted in more potent mTAAR1 agonists when compared to T1AM, as well as, lead compound SG2. Taking into account the high structure similarity of compounds **1** and **2** with T1AM and their high affinity for mTAAR1, we further investigated their ability to modulate glucose levels using both *in vitro* (human HepG2 cells) and *in vivo* (CD1-mice) models. Both

1
2
3 compounds demonstrated potent functional activity. In conclusion, this paper describes for the first
4
5 time the successful outcome of a hit-to-lead optimization process leading to the identification of
6
7 new tools to explore the physiological and pharmacological functions of TAAR1.
8

9 10 **Experimental Section**

11 12 *Chemistry.*

13
14
15 *General Material and Methods.* Melting points were determined on a Kofler hot-stage apparatus
16
17 and are uncorrected. Chemical shifts (δ) are reported in parts per million downfield from
18
19 tetramethylsilane and referenced from solvent references; coupling constants J are reported in hertz.
20
21 ^1H NMR and ^{13}C NMR spectra of all compounds were obtained with a Bruker TopSpin 3.2 400
22
23 MHz spectrometer. ^{13}C NMR spectra were fully decoupled. The following abbreviations are used:
24
25 singlet (s), doublet (d), triplet (t), double-doublet (dd), and multiplet (m). The elemental
26
27 compositions of the compounds agreed to within $\pm 0.4\%$ of the calculated values. Chromatographic
28
29 separation was performed on silica gel columns by flash (Kieselgel 40, 0.040–0.063 mm; Merck) or
30
31 gravity column (Kieselgel 60, 0.063–0.200 mm; Merck) chromatography. The $\geq 95\%$ purity of the
32
33 tested compounds was confirmed by combustion analysis. Reactions were followed by thin-layer
34
35 chromatography (TLC) on Merck aluminum silica gel (60 F₂₅₄) sheets that were visualized under a
36
37 UV lamp. The microwave-assisted procedures were carried out with a CEM Discover LabMate
38
39 microwave. Evaporation was performed in vacuo (rotating evaporator). Sodium sulfate was always
40
41 used as the drying agent. Commercially available chemicals were purchased from Sigma-Aldrich.
42
43
44
45
46

47
48 *4-(4-(2-Aminoethyl)-2-methylbenzyl)aniline (1).* To a suspension of LiAlH_4 (4.80 mL, 4.80 mmol)
49
50 in THF was added dropwise a solution of AlCl_3 (639.6 mg, 4.80 mmol) in THF (20 mL), and the
51
52 mixture was stirred for 5 min at r.t. A solution of **25** (142.0 mg, 0.53 mmol) in THF was added
53
54 dropwise, and the mixture was heated at reflux for 12 h. The mixture was cooled to 0°C with an ice
55
56 bath and added dropwise with water and then with 10% aqueous HCl. The mixture was extracted
57
58 with diethyl ether, and the aqueous layer was made alkaline with NaOH 2N and extracted with
59
60

1
2
3 CHCl₃. The organic phase was separated, washed with brine, dried, filtered, and concentrated. The
4
5 crude was purified by conversion in the corresponding hydrochloride salt. White solid. mp: 153-
6
7 157°C. (67% yield). ¹H NMR (CD₃OD): δ 2.21 (s, 3H, CH₃); 2.94 (t, 2H, *J* = 7.8 Hz; CH₂); 3.17 (t,
8
9 2H, *J* = 7.8 Hz; CH₂); 4.03 (s, 2H, CH₂); 7.07-7.15 (m, 3H, Ar); 7.29 (d, 2H, *J* = 8.4 Hz, Ar); 7.34
10
11 (d, 2H, *J* = 8.4 Hz, Ar) ppm. ¹³C NMR (CD₃OD): δ 143.44, 138.48, 138.37, 136.40, 131.89, 131.66,
12
13 131.35, 129.71, 127.54, 124.11, 41.99, 39.31, 34.09, 19.70 ppm. Anal. (C₁₆H₂₀N₂·HCl) C, H, N. %
14
15 Calcd: 79.96 (C); 8.39 (H); 11.66 (N). % Found: 79.88 (C); 8.55 (H); 11.42 (N).
16
17

18
19 *4-(4-(2-Aminoethyl)-2-methylbenzyl)phenol (2)*. Compound **2** was synthesized from **30** (109 mg,
20
21 0.46 mmol), LiAlH₄ (4.13 mL; 4.13 mmol) and AlCl₃ (550.8 mg, 4.13 mmol) in THF (18 mL)
22
23 following the same procedure described above for the preparation of **1**. Yellow solid. mp: 138-
24
25 140°C. (30% yield). ¹H NMR (CD₃OD): δ 2.18 (s, 3H, CH₃); 2.70 (t, 2H, *J* = 7.2 Hz; CH₂); 2.87 (t,
26
27 2H, *J* = 7.2 Hz; CH₂); 3.81 (s, 2H, CH₂); 6.64 (d, 2H, *J*=8.4 Hz, Ar); 6.84 (d, 2H, *J*=8.4 Hz, Ar);
28
29 6.94-7.03 (m, 3H, Ar) ppm. ¹³C NMR (CD₃OD): δ 146.29, 139.28, 137.93, 137.82, 131.69, 131.64,
30
31 131.10, 130.31, 127.21, 116.88, 54.79, 39.22, 38.30, 19.74 ppm. Anal. (C₁₆H₁₉NO) C, H, N. %
32
33 Calcd: 79.63 (C); 7.94 (H); 5.80 (N). % Found: 79.77 (C); 7.81 (H); 5.92 (N).
34
35
36

37
38 *N-(4-(4-(2-Aminoethyl)-2-methylbenzyl)phenyl)acetamide (3)*. To a solution of **1** (50 mg, 0.18
39
40 mmoli) in saturated NaHCO₃ solution (0.06 mL) was added Ac₂O (0.02 mL, 0.18 mmoli). The
41
42 mixture was left stirring at r.t. for 1 h, then, the mixture was dissolved in water and extracted with
43
44 CHCl₃. The organic layer was dried and concentrated under reduced pressure. The crude product
45
46 was purified by precipitation from MeOH/Et₂O to give **3**. White solid. mp: 132-136°C. (60% yield).
47
48 ¹H NMR (CD₃OD): δ 2.10 (s, 3H, CH₃); 2.20 (s, 3H, COCH₃); 2.92 (t, 2H, *J* = 7.7 Hz; CH₂); 3.15
49
50 (t, 2H, *J* = 7.7 Hz; CH₂); 3.91 (s, 2H, CH₂); 7.04 (d, 2H, *J* = 8.4 Hz, Ar); 7.05-7.11 (m, 3H, Ar);
51
52 7.34 (d, 2H, *J* = 8.4 Hz, Ar) ppm. ¹³C NMR (CD₃OD): δ 171.57, 139.37, 138.31, 137.77, 137.62,
53
54 135.91, 131.71, 131.52, 129.93, 127.33, 121.33, 42.01, 39.37, 34.09, 23.77, 19.74 ppm. Anal.
55
56
57
58
59
60

1
2
3 (C₁₆H₂₀N₂·HCl) C, H, N. % Calcd: 79.96 (C); 8.39 (H); 11.66 (N). % Found: 79.88 (C); 8.55 (H);
4
5 11.42 (N).
6

7
8 *4-(4-(2-Aminoethyl)-2,6-dimethylbenzyl)aniline (4)*. Compound **4** was synthesized from **26** (109
9
10 mg, 0.46 mmol), LiAlH₄ (4.13 mL; 4.13 mmol) and AlCl₃ (550.8 mg, 4.13 mmol) in THF (18 mL)
11
12 following the same procedure described above for the preparation of **1**. The crude was purified by
13
14 conversion in the corresponding hydrochloride salt. White solid, mp: 143-145°C. (68% yield). ¹H
15
16 NMR (CD₃OD): δ 2.22 (s, 6H, CH₃); 2.91 (t, 2H, *J* = 7.7 Hz; CH₂); 3.18 (t, 2H, *J* = 7.7 Hz; CH₂);
17
18 4.11 (s, 2H, CH₂); 7.01 (s, 2H, Ar); 7.17 (d, 2H, *J* = 8.6 Hz, Ar); 7.30 (d, 2H, *J* = 8.6 Hz, Ar) ppm.
19
20 ¹³C NMR (CD₃OD): δ 142.77, 138.85, 136.31, 136.14, 130.58, 129.68, 129.61, 124.13, 42.00,
21
22 34.98, 34.12, 20.23 ppm. Anal. (C₁₇H₂₂N₂·HCl) C, H, N. % Calcd: 80.27 (C); 8.72 (H); 11.01 (N).
23
24 % Found: 80.40 (C); 8.59 (H); 11.31 (N).
25
26
27

28
29 *4'-(2-Aminoethyl)-2'-methyl-[1,1'-biphenyl]-4-amine (5)*. Compound **5** was synthesized from **27**
30
31 (109 mg, 0.46 mmol), LiAlH₄ (4.13 mL; 4.13 mmol) and AlCl₃ (550.8 mg, 4.13 mmol) in THF (18
32
33 mL) following the same procedure described above for the preparation of **1**. The crude was purified
34
35 by conversion in the corresponding hydrochloride salt. White solid, mp: 165-167°C. (70% yield).
36
37 ¹H NMR (CD₃OD): δ 2.25 (s, 3H, CH₃); 2.99 (t, 2H, *J* = 7.8 Hz; CH₂); 3.22 (t, 2H, *J* = 7.8 Hz;
38
39 CH₂); 7.19 (d, 2H, *J* = 1.2 Hz, Ar); 7.25 (s, 1H, Ar); 7.51-7.46 (m, 4H, Ar) ppm. ¹³C NMR
40
41 (CD₃OD): δ 144.11, 140.55, 137.72, 137.02, 131.99, 131.21, 130.72, 127.48, 124.08, 41.92, 34.19,
42
43 20.43 ppm. Anal. (C₁₅H₁₈N₂·HCl) C, H, N. % Calcd: 79.61 (C); 8.02 (H); 12.38 (N). % Found:
44
45 80.00 (C); 8.10 (H); 12.22 (N).
46
47
48

49
50 *4'-(2-Aminoethyl)-2',6'-dimethyl-[1,1'-biphenyl]-4-amine (6)*. Compound **6** was synthesized from **28**
51
52 (109 mg, 0.46 mmol), LiAlH₄ (4.13 mL; 4.13 mmol) and AlCl₃ (550.8 mg, 4.13 mmol) in THF (18
53
54 mL) following the same procedure described above for the preparation of **1**. The crude was purified
55
56 by conversion to the corresponding hydrochloride salt. White solid, mp: 171-173°C. (73% yield).
57
58 ¹H NMR (CD₃OD): δ 2.00 (s, 3H, CH₃); 2.95 (t, 2H, *J* = 7.8 Hz; CH₂); 3.20 (t, 2H, *J* = 7.8 Hz;
59
60

1
2
3 CH₂); 7.07 (s, 2H, Ar); 7.30 (d, 2H, *J* = 8.4 Hz, Ar); 7.53 (d, 2H, *J* = 8.4 Hz, Ar) ppm. ¹³C NMR
4 (CD₃OD): δ 143.35, 140.41, 137.50, 137.27, 132.17, 130.72, 128.97, 124.52, 41.98, 34.21, 20.84
5
6 ppm. Anal. (C₁₆H₂₀N₂·HCl) C, H, N. % Calcd: 79.96 (C); 8.39 (H); 11.66 (N). % Found: 79.90 (C);
7
8 8.49 (H); 11.71 (N).
9

10
11
12 *4-(4-(2-Aminoethoxy)-2,6-dimethylbenzyl)aniline (7)*. Compound **7** was synthesized from **33** (186.2
13 mg, 0.66 mmol), LiAlH₄ (6.87 mL, 6.87 mmol) and AlCl₃ (915.6 mg, 6.87 mmol) in THF (25 mL)
14 following the same procedure described above for the preparation of **1**. The crude was purified by
15 conversion to the corresponding hydrochloride salt. White solid, mp: 156-158°C. (75% yield). ¹H
16 NMR (CD₃OD): δ 2.20 (s, 6H, CH₃); 3.37 (t, 2H, *J* = 5.0 Hz; CH₃); 4.07 (s, 2H, CH₂); 4.23 (t, 2H, *J*
17 = 5.0 Hz; CH₂); 6.76 (s, 2H, Ar); 7.15 (d, 2H, *J* = 8.4 Hz, Ar); 7.30 (d, 2H, *J* = 8.4 Hz, Ar) ppm. ¹³C
18 NMR (CD₃OD): δ 156.50, 141.82, 138.32, 129.24, 129.09, 128.16, 122.72, 114.04, 63.74, 39.07,
19 33.21, 19.10 ppm. Anal. (C₁₇H₂₂N₂O·HCl) C, H, N. % Calcd: 75.52 (C); 8.20 (H); 10.36 (N). %
20 Found: 75.60 (C); 8.07 (H); 10.40 (N).
21
22
23
24
25
26
27
28
29
30
31
32

33
34 *4-(4-(2-Aminoethyl)-2-methylbenzyl)-N-ethyl-2-isopropylaniline (8)*. Compound **8** was synthesized
35 from **37** (130.8 mg, 0.41 mmol), LiAlH₄ (1.84 mL, 1.84 mmol) and AlCl₃ (244.8 mg, 1.84 mmol) in
36 THF (17 mL) following the same procedure described above for the preparation of **1**. The crude
37 was purified by conversion in the corresponding hydrochloride salt. Brown solid, mp: 148-150°C.
38 (67% yield). ¹H NMR (CD₃OD): δ 1.28 (d, 6H, *J* = 6.8 Hz, CH₃); 1.37 (t, 3H, *J* = 7.2 Hz, CH₃);
39 2.22 (s, 3H, CH₃); 2.93 (t, 2H, *J* = 7.8 Hz, CH₂); 3.05-3.12 (m, 1H, CH); 3.16 (t, 2H, *J* = 7.8 Hz,
40 CH₂); 3.43 (q, 2H, *J* = 7.2 Hz, CH₂); 4.04 (s, 2H, CH₂); 7.06-7.14 (m, 4H, Ar); 7.32 (d, 1H, *J* = 8.4
41 Hz, Ar); 7.36 (d, 1H, *J* = 1.6 Hz, Ar) ppm. ¹³C NMR (CD₃OD): δ 144.42, 143.51, 138.47, 138.36,
42 136.41, 131.87, 131.56, 131.00, 129.52, 128.88, 127.54, 124.53, 49.90, 41.97, 39.45, 34.11, 28.70,
43 24.43, 19.72, 11.38 ppm. Anal. (C₂₁H₃₀N₂·HCl) C, H, N. % Calcd: 81.24 (C); 9.74 (H); 9.02 (N). %
44 Found: 81.36 (C); 9.65 (H); 9.11 (N).
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *N*-(4-(4-(2-Aminoethyl)-2-methylbenzyl)-2-isopropylphenyl)-*N*-ethylacetamide (**9**). Compound **9**
4 was synthesized from **8** (45.6 mg, 0.15 mmol), saturated NaHCO₃ solution (0.05 mL) and Ac₂O
5 (0.01 mL, 0.15 mmol) following the same procedure described above for the preparation of **3**. The
6 crude solid was collected by filtration and purified by conversion to the corresponding
7 hydrochloride salt. Yellow solid. (30% yield). ¹H NMR (CD₃OD): δ 1.08-1.20 (m, 9H, CH₃,
8 CH₃CH₂); 1.73 (s, 3H, COCH₃); 2.24 (s, 3H, CH₃); 2.90-2.96 (m, 3H, CH₂, CH); 3.02-3.07 (m, 1H,
9 CH₂); 3.17 (t, 2H, *J* = 7.4 Hz, CH₂); 4.02 (s, 2H, CH₂); 4.11-4.18 (m, 1H, CH₂); 6.97-7.05 (m, 2H,
10 Ar); 7.07-7.17 (m, 3H, Ar); 7.27 (s, 1H, Ar) ppm. ¹³C NMR (CD₃OD): δ 173.03, 146.93, 143.12,
11 138.91, 138.71, 138.36, 136.21, 131.81, 131.53, 130.08, 128.87, 128.30, 127.47, 45.29, 42.01,
12 39.67, 34.10, 28.83, 24.48, 24.13, 22.59, 19.77, 12.99 ppm. Anal. (C₂₃H₃₂N₂O·HCl) C, H, N. %
13 Calcd: 78.36 (C); 9.15 (H); 7.95 (N). % Found: 78.45 (C); 9.02 (H); 7.84 (N).
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 *N*-(4-(4-(2-Acetamidoethyl)-2-methylbenzyl)-2-isopropylphenyl)-*N*-ethylacetamide (**10**). Compound
29 **10** was synthesized from **8** (45.6 mg, 0.15 mmol), saturated NaHCO₃ solution (0.05 mL) and Ac₂O
30 (0.02 mL, 0.30 mmol) following the same procedure described above for the preparation of **3**. The
31 reaction mixture was evaporated and the crude dissolved in AcOEt and washed twice with water.
32 The evaporation of the organic collected phases gave final product as a white oil. (20% yield). ¹H
33 NMR (CD₃OD): δ 1.10-1.20 (m, 9H, CH₃, CH₃CH₂); 1.73 (s, 3H, COCH₃); 1.90 (s, 3H, COCH₃);
34 2.21 (s, 3H, CH₃); 2.74 (t, 2H, *J* = 7.4 Hz, CH₂); 2.90-2.96 (m, 1H, CH); 3.01-3.09 (m, 1H, CH₂);
35 3.37 (t, 2H, *J* = 7.4 Hz, CH₂); 4.00 (s, 2H, CH₂); 4.11-4.20 (m, 1H, CH₂); 6.99-7.09 (m, 5H, Ar);
36 7.25 (d, 1H, *J* = 1.2 Hz, Ar) ppm. ¹³C NMR (CD₃OD): δ 173.20, 173.08, 146.89, 143.36, 138.82,
37 138.68, 137.88, 137.74, 131.86, 131.13, 130.06, 128.84, 128.28, 127.48, 45.34, 42.13, 39.71, 36.09,
38 28.86, 24.50, 24.16, 22.60, 22.52, 19.78, 13.00 ppm. Anal. (C₂₅H₃₄N₂O₂) C, H, N. % Calcd: 76.10
39 (C); 8.69 (H); 7.10 (N). % Found: 76.25 (C); 8.53 (H); 7.21 (N).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 *N*-(4-(4-(2-Aminoethoxy)benzyl)phenyl)acetamide (**11**). Compound **11** was synthesized from SG1⁴
56 (49.7 mg, 0.20 mmol), saturated NaHCO₃ solution (0.07 mL) and Ac₂O (0.03 mL, 0.20 mmol)
57
58
59
60

1
2
3 following the same procedure described above for the preparation of **3**. The crude product was
4
5 purified by precipitation from MeOH/Et₂O to give **11**. White solid, mp: 132-136°C. (60% yield). ¹H
6
7 NMR (CD₃OD): δ 2.10 (s, 3H, COCH₃); 3.34 (t, 2H, *J* = 5.2 Hz; CH₂); 3.87 (s, 2H, CH₂); 4.19 (t,
8
9 2H, *J* = 5.2 Hz; CH₂); 6.92 (d, 2H, *J* = 8.8 Hz, Ar); 7.10-7.14 (m, 4H, Ar); 7.42 (d, 2H, *J* = 8.4 Hz,
10
11 Ar) ppm. ¹³C NMR (CD₃OD): δ 171.59, 157.83, 138.90, 137.82, 136.02, 130.93, 130.03, 121.40,
12
13 115.70, 65.29, 41.30, 40.39, 23.76 ppm. Anal. (C₁₇H₂₀N₂·HCl) C, H, N. % Calcd: 71.81 (C); 7.09
14
15 (H); 9.85 (N). % Found: 71.88 (C); 7.15 (H); 9.92 (N).
16
17
18

19 *N*-(2-(4-(4-Acetamidobenzyl)phenoxy)ethyl)acetamide (**12**). Compound **12** was synthesized from
20
21 **SG1** (47.0 mg, 0.19 mmol), saturated NaHCO₃ solution (0.14 mL) and Ac₂O (0.04 mL, 0.38 mmol)
22
23 following the same procedure described above for the preparation of **3**. The crude product was
24
25 purified by precipitation from MeOH/Acetone to give **12**. White solid, mp: 152-153°C. (60% yield).
26
27 ¹H NMR (DMSO): δ 2.00 (s, 3H, COCH₃); 2.08 (s, 3H, COCH₃); 3.34-3.38 (m, 2H, CH₂); 3.79 (s,
28
29 2H, CH₂); 3.91 (t, 2H, *J* = 5.6 Hz; CH₂); 6.84 (d, 2H, *J* = 8.8 Hz, Ar); 7.08-7.10 (m, 4H, Ar); 7.45
30
31 (d, 2H, *J* = 8.8 Hz, Ar); 8.06 (br t, NH₂); 9.83 (br s, NH) ppm. ¹³C NMR (CD₃OD): δ 173.59,
32
33 171.54, 158.58, 139.08, 137.78, 135.20, 130.82, 130.06, 121.39, 115.59, 67.53, 41.32, 40.26, 23.70,
34
35 22.46 ppm. Anal. (C₁₉H₂₂N₂O₃·HCl) C, H, N. % Calcd: 69.92 (C); 6.79 (H); 8.58 (N). % Found:
36
37 69.88 (C); 6.67 (H); 8.61 (N).
38
39
40
41

42 *N*-(4-(4-(2-Aminoethoxy)-2-methylbenzyl)phenyl)acetamide (**13**). Compound **13** was synthesized
43
44 from **SG2**⁴ (50.0 mg, 0.19 mmol), saturated NaHCO₃ solution (0.07 mL) and Ac₂O (0.02 mL, 0.19
45
46 mmoli) following the same procedure described above for the preparation of **3**. The crude product
47
48 was purified by precipitation from MeOH/Et₂O to give **13**. White solid, mp: 132-136°C. (60%
49
50 yield). ¹H NMR (CD₃OD): δ 2.10 (s, 3H, CH₃); 2.19 (s, 3H, COCH₃); 3.34 (t, 2H, *J* = 5.0 Hz; CH₂);
51
52 3.89 (s, 2H, CH₂); 4.20 (t, 2H, *J* = 5.0 Hz; CH₂); 6.77 (dd, 1H, *J* = 2.4, 8.4 Hz, Ar); 6.83 (d, 1H, *J* =
53
54 2.4 Hz, Ar); 7.03 (d, 2H, *J* = 8.4 Hz, Ar); 7.06 (d, 1H, *J* = 8.4 Hz, Ar); 7.41 (d, 2H, *J* = 8.4 Hz, Ar)
55
56 ppm. ¹³C NMR (CD₃OD): δ 171.58, 158.00, 139.19, 138.03, 137.67, 133.61, 132.00, 129.79,
57
58
59
60

1
2
3 121.32, 117.73, 112.83, 65.18, 40.41, 38.93, 23.79, 19.96 ppm. Anal. (C₁₈H₂₂N₂O₂·HCl) C, H, N.
4
5 % Calcd: 72.46 (C); 7.43 (H); 9.39 (N). % Found: 72.58 (C); 7.55 (H); 9.42 (N).
6
7

8 *N*-(2-(4-(4-Acetamidobenzyl)-3-methylphenoxy)ethyl)acetamide (**14**). Compound **14** was
9
10 synthesized from **SG2** (50.0 mg, 0.19 mmol), saturated NaHCO₃ solution (0.14 mL) and Ac₂O
11 (0.04 mL, 0.38 mmol) following the same procedure described above for the preparation of **3**. The
12 crude product was purified by precipitation from MeOH/Et₂O to give **14**. White solid, mp: 139-
13 141°C. (56% yield). ¹H NMR (CD₃OD): δ 1.95 (s, 3H, CH₃); 2.09 (s, 3H, COCH₃); 2.16 (s, 3H,
14 COCH₃); 3.53 (t, 2H, *J* = 5.6 Hz; CH₂); 3.87 (s, 2H, CH₂); 4.00 (t, 2H, *J* = 5.6 Hz; CH₂); 6.70 (dd,
15 1H, *J* = 2.4, 8.4 Hz, Ar); 6.74 (d, 1H, *J* = 2.4 Hz, Ar); 6.99-7.03 (m, 3H, Ar); 7.41 (d, 2H, *J* = 8.4
16 Hz, Ar) ppm. ¹³C NMR (CD₃OD): δ 173.60, 171.53, 158.78, 139.01, 138.27, 137.66, 132.85,
17 131.94, 129.83, 121.32, 117.66, 112.72, 67.42, 40.30, 38.98, 23.70, 22.47, 19.96 ppm. Anal.
18 (C₂₀H₂₄N₂O₃·HCl) C, H, N. % Calcd: 70.56 (C); 7.11 (H); 8.23 (N). % Found: 70.43 (C); 7.05 (H);
19 8.22 (N).
20
21
22
23
24
25
26
27
28
29
30
31
32

33 *N*-(4-(4-(2-(3-Carbamimidoylguanidino)ethoxy)-2-methylbenzyl)phenyl)acetamide (**15**). A mixture
34 of **13** (32.3 mg, 0.16 mmol) and dicyandiamide (13.5 mg, 0.16 mmol) was heated to 160 °C for 100
35 min. The reaction mixture initially melts, then re-solidifies. The reaction was cooled to room
36 temperature. The crude product was purified by crystallization from EtOH/Et₂O. Brown oil. (52%
37 yield). ¹H NMR (CD₃OD): δ 2.09 (s, 3H, CH₃); 2.15 (s, 3H, COCH₃); 3.60 (t, 2H, *J* = 5.2 Hz; CH₂);
38 3.86 (s, 2H, CH₂); 4.05 (t, 2H, *J* = 5.2 Hz; CH₂); 6.65-6.77 (m, 3H, Ar); 7.01 (d, 2H, *J* = 8.0 Hz,
39 Ar); 7.41 (d, 2H, *J* = 8.0 Hz, Ar) ppm. ¹³C NMR (CD₃OD): δ 171.60, 162.12, 161.01, 158.56,
40 139.04, 138.20, 137.63, 133.00, 131.94, 129.80, 121.34, 117.63, 112.73, 67.55, 42.15, 38.94, 23.74,
41 19.97 ppm. Anal. (C₂₀H₂₆N₆O₂) C, H, N. % Calcd: 62.81 (C); 6.85 (H); 21.97 (N). % Found: 62.95
42 (C); 6.73 (H); 21.76 (N).
43
44
45
46
47
48
49
50
51
52
53
54
55

56 *4*-(4-(2-(3-Carbamimidoylguanidino)ethoxy)-2-methylbenzyl)aniline (**16**). To a solution of the
57 amide **15** (0.57 mmol) in MeOH was added dropwise an aqueous solution of HCl 1N (0.2 mL); the
58
59
60

1
2
3 resulting solution was refluxed for 2h, then, after cooling, the solvent was evaporated. The residue
4
5 was purified by precipitation from EtOH/Et₂O. Brown oil. (60% yield). ¹H NMR (CD₃OD): δ 2.17
6
7 (s, 3H, CH₃); 3.79 (t, 2H, *J* = 4.8 Hz; CH₂); 3.99 (s, 2H, CH₂); 4.21 (t, 2H, *J* = 4.8 Hz; CH₂); 6.79
8
9 (dd, 1H, *J* = 2.6, 8.2, Ar); 6.83 (d, 1H, *J* = 2.6 Hz, Ar); 7.08 (d, 1H, *J* = 8.2 Hz, Ar), 7.26 (d, 2H, *J* =
10
11 8.6 Hz, Ar), 7.32 (d, 2H, *J* = 8.6 Hz, Ar) ppm. Anal. (C₁₈H₂₄N₆O) C, H, N. % Calcd: 63.51 (C);
12
13 7.11 (H); 24.69 (N). % Found: 63.72 (C); 7.05 (H); 24.53 (N).

14
15
16
17 *4-(Chloromethyl)-2-methyl-1-(4-nitrobenzyl)benzene (21)*. To a solution of alcohol **17** (96.0mg,
18
19 0.35 mmol) in CHCl₃ at 0°C was added SOCl₂ (3.50 mmol; 0.25 mL). The reaction mixture was
20
21 stirred for 2 h at room temperature, then, the solvent was evaporated. The residue was dissolved in
22
23 H₂O and alkalized with NaOH 1N; and the aqueous layer was extracted with DCM. The organic
24
25 phase was dried and the solvent was evaporated to give the corresponding crude **21**, which was used
26
27 without further purification. Yellow oil. (80% yield). ¹H NMR (CDCl₃): δ 2.22 (s, 3H, CH₃); 4.08
28
29 (s, 2H, CH₂); 4.57 (s, 2H, CH₂); 7.09 (d, 1H, *J* = 7.6 Hz, Ar); 7.19-7.24 (m, 2H, Ar); 7.26 (d, 2H,
30
31 *J* = 7.4 Hz, Ar); 8.13 (d, 2H, *J* = 7.4 Hz, Ar) ppm. Anal. (C₁₅H₁₄ClNO₂) C, H, N. % Calcd: 65.34 (C);
32
33 5.12 (H); 5.08 (N). % Found: 65.52 (C); 5.03 (H); 5.24 (N).

34
35
36
37
38 *5-(Chloromethyl)-1,3-dimethyl-2-(4-nitrobenzyl)benzene (22)*. Compound **22** was synthesized from
39
40 **18** (224 mg, 0.87 mmol) and SOCl₂ (0.63 mL, 8.70 mmol) in CHCl₃ (25 mL) following the same
41
42 procedure described above for the preparation of **21**. The crude was used without further
43
44 purification. White oil. (76% yield). ¹H NMR (CDCl₃): δ 2.22 (s, 6H, CH₃); 4.14 (s, 2H, CH₂); 4.56
45
46 (s, 2H, CH₂); 7.13 (s, 2H, Ar); 7.15 (d, 2H, *J* = 8.5 Hz Ar); 8.11 (d, 2H, *J* = 8.5 Hz Ar) ppm. Anal.
47
48 (C₁₆H₁₆ClNO₂) C, H, N. % Calcd: 66.32 (C); 5.57 (H); 4.83 (N). % Found: 66.49 (C); 5.59 (H);
49
50 5.11 (N).

51
52
53
54 *4-(Chloromethyl)-2-methyl-4'-nitro-1,1'-biphenyl (23)*. Compound **23** was synthesized from **19** (125
55
56 mg, 0.51 mmol) and SOCl₂ (0.37 mL, 5.10 mmol) in CHCl₃ (25 mL) following the same procedure
57
58 described above for the preparation of **21**. The crude was used without further purification. Yellow
59
60

oil. (83% yield). $^1\text{H NMR}$ (CDCl_3): δ 2.27 (s, 3H, CH_3); 4.61 (s, 2H, CH_2); 7.21 (d, 2H, $J = 7.8$ Hz, Ar); 7.31 (d, 1H, $J = 7.8$ Hz, Ar); 7.34 (s, 1H, Ar); 7.48 (d, 2H, $J = 8.8$ Hz, Ar); 8.29 (d, 2H, $J = 8.8$ Hz Ar) ppm. Anal. ($\text{C}_{14}\text{H}_{12}\text{ClNO}_2$) C, H, N. % Calcd: 64.25 (C); 4.62 (H); 5.35 (N). % Found: 63.91 (C); 4.68 (H); 5.40 (N).

4-(Chloromethyl)-2,6-dimethyl-4'-nitro-1,1'-biphenyl (**24**). Compound **24** was synthesized from **20** (134 mg, 0.52 mmol) and SOCl_2 (0.38 mL, 5.20 mmol) in CHCl_3 (25 mL) following the same procedure described above for the preparation of **21**. The crude was used without further purification. White oil. (81% yield). $^1\text{H NMR}$ (CDCl_3): δ 2.02 (s, 6H, CH_3); 4.58 (s, 2H, CH_2); 7.17 (s, 2H, Ar); 7.33 (d, 2H, $J = 8.8$ Hz Ar); 8.31 (d, 2H, $J = 8.8$ Hz Ar) ppm. Anal. ($\text{C}_{15}\text{H}_{14}\text{ClNO}_2$) C, H, N. % Calcd: 65.34 (C); 5.12 (H); 5.08 (N). % Found: 65.40 (C); 5.59 (H); 5.31 (N).

2-(4-(4-Aminobenzyl)-3-methylphenyl)acetonitrile (**25**). To a solution of chloro compound **21** (78.0 mg; 0.46 mmol) in CH_3CN (0.62 mL) was added NaCN (45.1 mg; 0.92 mmol) in H_2O (0.21 mL). The mixture was submitted to microwave irradiation (150W, 100°C , 20 min). After cooling the solution was extracted with DCM. Organic phase was dried and evaporated to dryness. Yellow oil. (86% yield). $^1\text{H NMR}$ (CDCl_3): δ 2.22 (s, 3H, CH_3); 3.71 (s, 2H, CH_2); 4.07 (s, 2H, CH_2); 7.09-7.18 (m, 3H, Ar); 7.25 (d, 2H, $J = 8.6$ Hz, Ar); 8.13 (d, 2H, $J = 8.6$ Hz, Ar) ppm. Anal. ($\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_2$) C, H, N. % Calcd: 72.16 (C); 5.30 (H); 10.52 (N). % Found: 72.34 (C); 5.02 (H); 10.65 (N).

2-(3,5-Dimethyl-4-(4-nitrobenzyl)phenyl)acetonitrile (**26**). Compound **26** was synthesized from **22** (193 mg, 0.70 mmol) and NaCN (68.5 mg, 1.40 mmol) in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (0.94 mL/0.32 mL) following the same procedure described above for the preparation of **25**. The crude was used without further purification. White oil. (70% yield). $^1\text{H NMR}$ (CDCl_3): δ 2.22 (s, 6H, CH_3); 3.70 (s, 2H, CH_2); 4.13 (s, 2H, CH_2); 7.06 (s, 2H, Ar); 7.14 (d, 2H, $J = 8.6$ Hz Ar); 8.11 (d, 2H, $J = 8.6$ Hz Ar) ppm. Anal. ($\text{C}_{17}\text{H}_{16}\text{N}_2\text{O}_2$) C, H, N. % Calcd: 72.84 (C); 5.75 (H); 9.99 (N). % Found: 72.60 (C); 5.60 (H); 10.02 (N).

1
2
3 2-(2-Methyl-4'-nitro-[1,1'-biphenyl]-4-yl)acetonitrile (**27**). Compound **27** was synthesized from **23**
4 (95.0 mg, 0.61 mmol) and NaCN (59.8 mg, 1.22 mmol) in CH₃CN/H₂O (0.82 mL/0.30 mL)
5 following the same procedure described above for the preparation of **25**. The crude was used
6 without further purification. Yellow oil. (86% yield). ¹H NMR (CDCl₃): δ 2.28 (s, 3H, CH₃); 3.78
7 (s, 2H, CH₂); 7.21-7.31 (m, 3H, Ar); 7.47 (d, 2H, *J* = 7.0 Hz, Ar); 8.29 (d, 2H, *J* = 7.0 Hz Ar) ppm.
8
9 Anal. (C₁₅H₁₂N₂O₂) C, H, N. % Calcd: 71.42 (C); 4.79 (H); 11.10 (N). % Found: 71.41 (C); 4.80
10 (H); 11.15 (N).
11
12
13
14
15
16
17

18
19 2-(2,6-Dimethyl-4'-nitro-[1,1'-biphenyl]-4-yl)acetonitrile (**28**). Compound **28** was synthesized from
20 **24** (101 mg, 0.36 mmol) and NaCN (35.3 mg, 0.72 mmol) in CH₃CN/H₂O (0.49 mL/0.16 mL)
21 following the same procedure described above for the preparation of **25**. The crude was used
22 without further purification. Pale yellow oil. (81% yield). ¹H NMR (CDCl₃): δ 2.02 (s, 6H, CH₃);
23 3.73 (s, 2H, CH₂); 7.10 (s, 2H, Ar); 7.32 (d, 2H, *J* = 8.6 Hz Ar); 8.31 (d, 2H, *J* = 8.6 Hz Ar) ppm.
24
25 Anal. (C₁₆H₁₄N₂O₂) C, H, N. % Calcd: 72.16 (C); 5.30 (H); 10.52 (N). % Found: 72.40 (C); 5.59
26 (H); 10.31 (N).
27
28
29
30
31
32
33
34

35 2-(4-(4-Aminobenzyl)-3-methylphenyl)acetonitrile (**29**). A solution of **25** (136 mg, 0.5 mmol) in
36 AcOH (0.5 mL) was hydrogenated in the presence of 10% Pd-C (28.3 mg), for 12 h. Then the
37 catalyst was filtered off, and the solvent was removed to dryness to give a crude product that was
38 used in the subsequent step without any further purification. Yellow oil. (80% yield). ¹H NMR
39 (CDCl₃) δ: 2.24 (s, 3H, CH₃); 3.68 (s, 2H, CH₂); 3.85 (s, 2H, CH₂); 6.61 (d, 2H, *J* = 8.4 Hz, Ar);
40 6.88 (d, 2H, *J* = 8.4 Hz, Ar); 7.06-7.12 (m, 3H, Ar) ppm. Anal. (C₁₆H₁₆N₂) C, H, N. % Calcd: 81.32
41 (C); 6.82 (H); 11.85 (N). % Found: 81.45 (C); 6.73 (H); 11.98 (N).
42
43
44
45
46
47
48
49
50

51 2-(4-(4-Hydroxybenzyl)-3-methylphenyl)acetonitrile (**30**). To a mixture of aniline derivative **29**
52 (63.0 mg, 0.26 mmol) in H₂O was added dropwise with H₂SO₄conc (0.06 mL) and the mixture was
53 stirred at room temperature for 20 min. Then a solution NaNO₂ (17.9 mg, 0.26 mmol) in H₂O (0.19
54 ml) was added dropwise to the reaction mixture. The resulting solution was stirred for 1 h at 100°C.
55
56
57
58
59
60

1
2
3 The mixture was cooled to room temperature, and the residue diluted with AcOEt and washed with
4 brine. The collected organic layers were dried and evaporated to give compound **30** that was
5 directly used in the next step. Yellow oil. (60% yield). $^1\text{H NMR}$ (CDCl_3) δ : 2.22 (s, 3H, CH_3); 3.84
6 (s, 2H, CH_2); 4.06 (s, 2H, CH_2); 7.14-7.19 (m, 3H, Ar); 7.27-7.33 (m, 4H, Ar) ppm. Anal.
7 ($\text{C}_{16}\text{H}_{15}\text{NO}$) C, H, N. % Calcd: 80.98 (C); 6.37 (H); 5.90 (N). % Found: 80.72 (C); 6.51 (H); 5.84
8 (N).
9
10
11
12
13
14
15

16
17 *5-Methoxy-1,3-dimethyl-2-(4-nitrobenzyl)benzene (31)*. To a solution of 4-methoxy-2,6-
18 dimethylphenylboronic acid (473 mg, 2.63 mmol) in acetone/ H_2O 1:1 (4 mL) was added, under
19 nitrogen flux, *p*-nitrobenzyl bromide (569 mg, 2.63 mmol), K_2CO_3 (1.24 g, 6.58 mmol) and a
20 catalytic amount of PdCl_2 . The resulting mixture was stirred at r.t. for 62 h in a sealed vial. The
21 crude mixture was evaporated and then diluted with water and extracted with Et_2O . The organic
22 phase was dried over sodium sulfate and concentrated under vacuum. The crude residue was
23 purified by column chromatography Petroleum Ether /AcOEt (99:1), affording **31**. Yellow oil. (49%
24 yield). $^1\text{H NMR}$ (CDCl_3): δ 2.19 (s, 6H, CH_3); 3.80 (s, 3H, OCH_3); 4.08 (s, 2H, CH_2); 6.65 (s, 2H,
25 Ar); 7.16 (d, 2H, $J = 8.8$ Hz, Ar); 8.09 (d, 2H, $J = 8.8$ Hz, Ar) ppm. Anal. ($\text{C}_{16}\text{H}_{17}\text{NO}_3$) C, H, N. %
26 Calcd: 70.83 (C); 6.32 (H); 5.16 (N). % Found: 71.01 (C); 6.12 (H); 5.00 (N).
27
28
29
30
31
32
33
34
35
36
37
38
39

40 *3,5-Dimethyl-4-(4-nitrobenzyl)phenol (32)*. A solution of **31** (0.39 mmol) in anhydrous DCM
41 (1.5 mL) was cooled to -78 °C and treated dropwise with a solution of BBR_3 in DCM (3.94 mL,
42 1.24 mmol); the resulting solution was stirred at the same temperature for 5 min and at 0 °C for 1 h.
43 The mixture was then diluted with water and extracted with DCM. Organic phase was dried and
44 evaporated to give the product **32**. Pale yellow oil. (64% yield). $^1\text{H NMR}$ (CDCl_3): δ 2.16 (s, 6H,
45 CH_3); 4.06 (s, 2H, CH_2); 6.59 (s, 2H, Ar) 7.16 (d, 2H, $J=8.8$ Hz, Ar); 8.10 (d, 2H, $J=8.8$ Hz, Ar)
46 ppm. Anal. ($\text{C}_{15}\text{H}_{15}\text{NO}_3$) C, H, N. % Calcd: 70.02 (C); 5.88 (H); 5.44 (N). % Found: 70.00 (C); 5.91
47 (H); 5.63 (N).
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *2-(3,5-Dimethyl-4-(4-nitrobenzyl)phenoxy)acetonitrile (33)*. To a mixture of cesium carbonate (725
4 mg, 2.22 mmol) and phenol derivative **32** (0.44 mmol) in 50 mL of DMF was added BrCH₂CN
5 (0.03 mL, 0.44 mmol). The reaction mixture was stirred for 30 min at r.t., poured into 100mL of
6 cold HCl 1N and extracted with AcOEt. The organic phase was dried and evaporated to give the
7 product **33**. White oil. (98% yield). ¹H NMR (CDCl₃): δ 2.22 (s, 6H, CH₃); 4.10 (s, 2H, CH₂); 4.77
8 (s, 2H, CH₂CN); 6.73 (s, 2H, Ar); 7.15 (d, 2H, *J* = 8.8, Ar); 8.11 (d, 2H, *J* = 8.8 Hz, Ar) ppm. Anal.
9 (C₁₇H₁₆N₂O₃) C, H, N. % Calcd: 68.91 (C); 5.44 (H); 9.45 (N). % Found: 68.65 (C); 5.52 (H); 9.65
10 (N)
11
12
13
14
15
16
17
18
19

20
21 *N-(4-(4-(Hydroxymethyl)-2-methylbenzyl)-2-isopropylphenyl)acetamide (35)*. To a solution of
22 trifluoro(4-(hydroxymethyl)-2-methylphenyl)borane salt (509 mg; 2.38 mmol) in dioxane/H₂O 9:1
23 (4 mL) was added, under nitrogen atmosphere, N-(4-(bromomethyl)-2-isopropylphenyl)acetamide
24 **34** (513 mg; 2.38 mmol), cesium carbonate (2.30 g; 7.1 mmol) and PdCl₂dppf (34.8 mg; 0.05
25 mmol). The resulting mixture was stirred at 95°C for 24 h in a sealed vial. The crude mixture was
26 evaporated and then diluted with water and extracted with DCM. The organic phase was dried over
27 sodium sulfate and the solvent removed. The crude product was chromatographed on a silica gel
28 column, eluting with Petroleum Ether/AcOEt (9:1). Brown oil. (50% yield). ¹H NMR (CDCl₃): δ δ
29 1.20 (d, 6H, *J* = 6.8 Hz, CH₃); 2.19 (s, 3H, CH₃); 2.26 (s, 3H, CH₃CO); 2.96-3.03 (m, 1H, CH);
30 3.94 (s, 2H, CH₂); 4.65 (s, 2H, CH₂); 6.90 (dd, 1H, *J* = 1.6, 8.0 Hz, Ar), 7.05-7.09 (m, 2H, Ar); 7.13
31 (d, 1H, *J* = 8.0 Hz, Ar); 7.17 (s, 1H, Ar); 7.47 (d, 1H, *J* = 8.0 Hz) ppm. Anal. (C₂₀H₂₅NO₂) C, H, N.
32 % Calcd: 77.14 (C); 8.09 (H); 4.50 (N). % Found: 77.39 (C); 8.21 (H); 4.64 (N).
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 *N-(4-(4-Chloromethyl)-2-methylbenzyl)-2-isopropylphenyl)acetamide (36)*. Compound **36** was
50 synthesized from **35** (208 mg, 0.67 mmol) and SOCl₂ (0.49 mL, 6.70 mmol) in CHCl₃ (25 mL)
51 following the same procedure described above for the preparation of **21**. The crude was used
52 without any further purification. Brown oil. (70% yield). ¹H NMR (CDCl₃): δ 1.20 (d, 6H, *J* = 6.8
53 Hz, CH₃); 2.20 (s, 3H, CH₃); 2.26 (s, 3H, CH₃CO); 2.96-3.03 (m, 1H, CH); 3.93 (s, 2H, CH₂); 4.57
54
55
56
57
58
59
60

(s, 2H, CH₂); 6.90 (d, 1H, *J* = 6.0 Hz, Ar), 7.02-7.09 (m, 2H, Ar); 7.15 (d, 1H, *J* = 8.4 Hz, Ar); 7.19 (s, 1H, Ar); 7.48 (d, 1H, *J* = 8.4 Hz) ppm. Anal. (C₂₀H₂₄ClNO) C, H, N. % Calcd: 72.82 (C); 7.33 (H); 4.25 (N). % Found: 72.61 (C); 7.42 (H); 4.13 (N).

N-(4-(4-(Cyanomethyl)-2-methylbenzyl)-2-isopropylphenyl)acetamide (**37**). Compound **37** was synthesized from **36** (152 mg, 0.46 mmol) and NaCN (45.3 mg, 0.92 mmol) in CH₃CN/H₂O (1.86 mL/0.63 mL) following the same procedure described above for the preparation of **25**. The crude was used without any further purification. Yellow oil. (88% yield). ¹H NMR (CDCl₃): δ 1.20 (d, 6H, *J* = 6.8 Hz, CH₃); 2.19 (s, 3H, CH₃); 2.25 (s, 3H, CH₃CO); 2.96-3.03 (m, 1H, CH); 3.70 (s, 2H, CH₂); 3.93 (s, 2H, CH₂); 6.89 (d, 1H, *J* = 8.0 Hz, Ar), 7.03-7.09 (m, 3H, Ar); 7.12 (s, 1H, Ar); 7.48 (d, 1H, *J* = 8.0 Hz, Ar) ppm. Anal. (C₂₁H₂₄N₂O) C, H, N. % Calcd: 78.71 (C); 7.55 (H); 8.74 (N). % Found: 78.85 (C); 7.41 (H); 8.90 (N).

Molecular modeling

All compounds were built, parameterised (Gasteiger-Huckel method) and energy minimised within MOE using MMFF94 forcefield [MOE: Chemical Computing Group Inc. Montreal. H3A2R7 Canada. <http://www.chemcomp.com>]. For all compounds, the protonated form was considered for the *in silico* analyses.

Docking studies. Docking studies were performed starting from the in-house homology model of the murine TAAR1 receptor⁴, built on the X-ray structure of the human β₂-adrenoreceptor (PDB ID: 3PDS)¹⁷, following a protocol we previously discussed exploring the binding mode of other TAAR1 ligands^{8, 9}. Briefly, the most promising compounds were docked into the putative ligand binding site by means of the Surflex docking module implemented in Sybyl-X1.0¹⁸.

Surflex-Dock uses an empirically derived scoring function based on the binding affinities of X-ray protein-ligand complexes. The final total score listed by Surflex-Dock is a weighted sum of non-linear functions involving van der Waals surface distances between the appropriate pairs of exposed

1
2
3 protein and ligand atoms, including hydrophobic, polar, repulsive, entropic and solvation and crash
4
5 terms represented in terms of a total score conferred to any calculated conformer. Then, for all the
6
7 compounds, the best docking geometries (selected on the basis of the SurFlex scoring functions)
8
9 were refined by ligand/receptor complex energy minimization (CHARMM27) by means of the
10
11 MOE software.
12

13 **In vitro Biological studies**

14
15
16
17 **Reagents.** All cell culture reagents and buffers were from Invitrogen (Carlsbad, CA) and Sigma (St.
18
19 Louis, MO). Coelenterazine *h* was purchased from Promega (Madison, WI). Plasmid containing the
20
21 cDNA for the mTAAR1 were generously donated from Hoffman-La Roche. EPAC cAMP BRET
22
23 sensor was produced as described¹⁹.
24

25 **Cell Culture and BRET experiment**

26
27
28
29 Human embryonic kidney 293 cells (HEK293T) were maintained in Dulbecco's Modified Eagle's
30
31 medium supplemented with 10% (vol/vol) of FBS, 2 mM l-glutamine and 0.05 mg/ml of
32
33 gentamicin at 37°C in a humidified atmosphere at 95% air and 5% CO₂. Transient transfections
34
35 were performed 24 h after cells seeding using lipofectamine 2000 protocol (Invitrogen). 5µg of
36
37 mTAAR1 and 4µg of EPAC for each milliliter of transfection solution were used for the
38
39 experiments. For the BRET experiments the cells were plated, 6 hours after transfection, in poly-D-
40
41 lysine coated 96-well microplates at a density of 70,000 cells per well in phenol red free Minimum
42
43 Essential Medium containing 2% of FBS, 10 mM Hepes, 2 mM L-glutamine. The cells were then
44
45 cultured for an additional 24 h. BRET experiment was conducted as already described⁷. Briefly, for
46
47 time course experiments, the plate was read immediately after the addition of the agonist and for
48
49 approximately 20 minutes. All the compounds were tested for screening at the initial concentration
50
51 of 10 µM. Then, for active compounds, a dose response was performed, in order to calculate the
52
53 EC₅₀ values. All the experiments were conducted in presence of the phosphodiesterase inhibitor
54
55 IBMX (Sigma) at the final concentration of 200 µM. Readings were collected using a Tecan Infinite
56
57
58
59
60

1
2
3 instrument that allows the sequential integration of the signals detected in the 465 to 505nm and 515
4
5 to 555 nm windows using filters with the appropriate band pass and by using iControl software. The
6
7 acceptor/donor ratio was calculated as previously described²⁰. Curve was fitted using a non-linear
8
9 regression and one site specific binding with GraphPad Prism 5. Data are representative of 4-5
10
11 independent experiments and are expressed as means±SEM.
12

13
14 **Cell culture and glucose production evaluation.** Human hepatocellular carcinoma cells (HepG2),
15
16 obtained from American Type Culture Collection (Manassas, VA, USA), were cultured in DMEM
17
18 supplemented with 10% (v/v) fetal bovine serum, 1 mM pyruvate, 100 U/ml penicillin, and 100
19
20 mg/ml streptomycin at 37°C in a humidified atmosphere containing 5% CO₂ and sub-cultured
21
22 before confluence. To assess glucose release, HepG2 were seeded into six-well plates (5x10⁵
23
24 cells/well) and grown to 80% of confluence with standard medium. As previously described⁹,
25
26 before treatment cells were washed twice with PBS and then exposed for 4 h to test compounds (1
27
28 and 2) (0.1, 1 and 10 µM) in 1 ml DMEM base, glucose- and phenol red-free, containing 100 U/ml
29
30 penicillin, 100 mg/ml streptomycin, and 4 mM L-glutamine, supplemented with 2 mM sodium
31
32 pyruvate and 20 mM sodium lactate (glucose production buffer) at 37 °C in 5% CO₂. Exogenous
33
34 T1AM (0.1, 1 and 10 µM) was used as control. Control cells were incubated with supplemented
35
36 DMEM containing DMSO (1–2 µl/well). Cell culture medium was then collected and glucose
37
38 concentration was measured with a colorimetric glucose assay kit (GAHK-20, Sigma-Aldrich),
39
40 following manufacturer's instruction. Glucose concentrations were referred to the total protein
41
42 content (Bradford 1976) of whole HepG2 lysates. Results are expressed as the mean ± S.E.M.
43
44 Differences between groups were analyzed by ANOVA. The threshold of statistical significance
45
46 was set at P<0.05. GraphPad Prism version 6.0 for Windows (GraphPad Software, San Diego, CA,
47
48 USA) was used for data processing and statistical analysis.
49
50
51
52
53
54

55
56 **Measurement of plasma glycaemia**
57
58
59
60

1
2
3 This investigation and animal use procedure complied with the National Institutes of Health Guide
4 for the Care and Use of Laboratory Animals (NIH Publications no. 80-23, revised 1996) and were
5 approved by the Animal Care Committee of the Department of Pharmacology, University of
6 Florence, in compliance with the European Communities Council Directive of 24 November 1986
7 (86/609/EEC). All efforts were made to minimize the number of animals used and their suffering.

8
9
10
11
12
13
14 Glycaemia was monitored in blood collected from the tail vein of 4 h fasted male mice (CD1 strain,
15 20-30g, from Envigo, Italy), who had received **1** and **2** (1.32, 4 and 11 $\mu\text{g}\cdot\text{kg}^{-1}$ i.p.) or saline (i.p.)
16 (n = 8 in each group). Glycaemia was evaluated by a glucrefractometer 15 min after the i.p.
17 injections, as described⁴. Data are expressed as mean \pm SEM of independent experiments. Statistical
18 analysis was performed by oneway ANOVA, followed by Student–Newman–Keuls multiple
19 comparison post hoc test; the threshold of statistical significance was set at $P < 0.05$. Data analysis
20 was performed by GraphPad Prism 6.0 statistical program (GraphPad software, San Diego, CA,
21 USA). The acceptor/donor ratio was calculated as previously described²⁰. Curve was fitted using a
22 non-linear regression and one site specific binding with GraphPad Prism 5. Data are representative
23 of 4-5 independent experiments and are expressed as means \pm SEM.

24 25 26 27 28 29 30 31 32 33 34 35 36 **ASSOCIATED CONTENT**

37
38
39 **Supporting Information.** Synthetic pathways and procedures for the preparation of compound **34**
40 and derivatives **11-16**. ¹H NMR and ¹³C NMR spectra of final compounds. The Supporting
41 Information is available free of charge on the ACS Publications website
42
43
44

45 46 47 **AUTHOR INFORMATION**

48
49 Corresponding Authors: *For G.C.: phone, +39 050 2218677; E-mail, g.chiellini@bm.med.unipi.it.
50
51 *For S.R.: phone, +39 050 2219582; E-mail, simona.rapposelli@farm.unipi.it.
52
53

54 55 56 **Notes**

57 The authors declare no competing financial interest.
58
59
60

ACKNOWLEDGMENTS

We thank Prof. Anna Maria Raspolli Galletti from the Department of Chemistry, University of Pisa (Italy) for allowing us to perform microwave-assisted reactions in her lab, and Prof. Thomas Scanlan for supplying us T1AM. This work was supported by a local grant from the University of Pisa (to G.C. and S.R.) and by the Russian Science Foundation (project N14-25-00065) (to R.R.G.)

ABBREVIATION USED

T1AM, 3-iodothyronamine; T0AM, thyronamine; β -PEA, β -phenylethylamine; mTAAR1, murine Trace-Amine Associated-Receptor 1;

References

1. Scanlan, T. S.; Suchland, K. L.; Hart, M. E.; Chiellini, G.; Huang, Y.; Kruzich, P. J.; Frascarelli, S.; Crossley, D. A.; Bunzow, J. R.; Ronca-Testoni, S. 3-Iodothyronamine is an endogenous and rapid-acting derivative of thyroid hormone. *Nat. Med.* **2004**, *10*, 638-642.
2. Scanlan, T. S. 3-Iodothyronamine (T(1)AM): a new player on the thyroid endocrine team? *Endocrinology* **2009**, *150*, 1108-1111.
3. Braulke, L. J.; Klingenspor, M.; DeBarber, A.; Tobias, S. C.; Grandy, D. K.; Scanlan, T. S.; Heldmaier, G. 3-Iodothyronamine: a novel hormone controlling the balance between glucose and lipid utilisation. *J. Comp. Physiol. B* **2008**, *178*, 167-177.
4. Chiellini, G.; Nesi, G.; Digiaco, M.; Malvasi, R.; Espinoza, S.; Sabatini, M.; Frascarelli, S.; Laurino, A.; Cichero, E.; Macchia, M.; Gainetdinov, R. R.; Fossa, P.; Raimondi, L.; Zucchi, R.; Rapposelli, S. Design, synthesis, and evaluation of thyronamine analogues as novel potent mouse Trace Amine Associated Receptor 1 (mTAAR1) Agonists. *J. Med. Chem.* **2015**, *58*, 5096-5107.
5. Chiellini, G.; Rapposelli, S.; Zucchi, R. Synthetic analogues of 3-iodothyronamine (t1am) and uses thereof. WO2015151068A1, 2015.

- 1
2
3 6. Hoefig, C. Thyroid hormone metabolites in cardiovascular health and disease. *A Symposium*
4 *on the Occasion of the Centennial Anniversary of Thyroxine Discovery yroid hormone metabolites*
5 *in cardiovascular health and disease*. Delphi, Greece, June 11, **2015**.
6
7
8
9
10 7. Espinoza, S.; Masri, B.; Salahpour, A.; Gainetdinov, R. R. BRET approaches to characterize
11 dopamine and TAAR1 receptor pharmacology and signaling. *Methods Mol. Biol.* **2013**, *964*, 107-
12 122.
13
14
15
16 8. Cichero, E.; Espinoza, S.; Gainetdinov, R. R.; Brasili, L.; Fossa, P. Insights into the structure
17 and pharmacology of the human Trace Amine-Associated Receptor 1 (hTAAR1): homology
18 modelling and docking studies. *Chem. Biol. Drug Des.* **2013**, *81*, 509-516.
19
20
21
22
23 9. Cichero, E.; Espinoza, S.; Franchini, S.; Guariento, S.; Brasili, L.; Gainetdinov, R. R.; Fossa,
24 P. Further insights into the pharmacology of the human Trace Amine-Associated Receptors:
25 discovery of novel ligands for TAAR1 by a virtual screening approach. *Chem. Biol. Drug Des.*
26 **2014**, *84*, 712-720.
27
28
29
30
31
32 10. Cichero, E.; Espinoza, S.; Tonelli, M.; Franchini, S.; Gerasimov, A. S.; Sorbi, C.;
33 Gainetdinov, R. R.; Brasili, L.; Fossa, P. A homology modelling-driven study leading to the
34 discovery of the first mouse trace amine-associated receptor 5 (TAAR5) antagonists.
35 *MedChemComm* **2016**, *7*, 353-364.
36
37
38
39
40
41 11. Wainscott, D. B.; Little, S. P.; Yin, T.; Tu, Y.; Rocco, V. P.; He, J. X.; Nelson, D. L.
42 Pharmacologic characterization of the cloned human trace amine-associated receptor1 (TAAR1)
43 and evidence for species differences with the rat TAAR1. *J. Pharmacol. Exp. Ther.* **2007**, *320*, 475-
44 485.
45
46
47
48
49
50 12. Reese, E. A.; Norimatsu, Y.; Grandy, M. S.; Suchland, K. L.; Bunzow, J. R.; Grandy, D. K.
51 Exploring the determinants of trace amine-associated receptor 1's functional selectivity for the
52 stereoisomers of amphetamine and methamphetamine. *J. Med. Chem.* **2014**, *57*, 378-390.
53
54
55
56
57
58
59
60

- 1
2
3 13. Regard, J. B.; Kataoka, H.; Cano, D. A.; Camerer, E.; Yin, L.; Zheng, Y.-W.; Scanlan, T. S.;
4
5 Hebrok, M.; Coughlin, S. R. Probing cell type-specific functions of Gi in vivo identifies GPCR
6
7 regulators of insulin secretion. *J. Clin. Invest.* **2007**, *117*, 4034-4043.
8
9
10 14. Klieverik, L. P.; Foppen, E.; Ackermans, M. T.; Serlie, M. J.; Sauerwein, H. P.; Scanlan, T.
11
12 S.; Grandy, D. K.; Fliers, E.; Kalsbeek, A. Central effects of thyronamines on glucose metabolism
13
14 in rats. *J. Endocrinol.* **2009**, *201*, 377-386.
15
16 15. Manni, M. E.; De Siena, G.; Saba, A.; Marchini, M.; Dicembrini, I.; Bigagli, E.; Cinci, L.;
17
18 Lodovici, M.; Chiellini, G.; Zucchi, R. 3-Iodothyronamine: a modulator of the hypothalamus-
19
20 pancreas-thyroid axes in mice. *Br. J. Pharmacol.* **2012**, *166*, 650-658.
21
22 16. Ghelardoni, S.; Chiellini, G.; Frascarelli, S.; Saba, A.; Zucchi, R. Uptake and metabolic
23
24 effects of 3-iodothyronamine in hepatocytes. *J. Endocrinol.* **2014**, *221*, 101-110.
25
26 17. Rosenbaum, D. M.; Zhang, C.; Lyons, J. A.; Holl, R.; Aragao, D.; Arlow, D. H.; Rasmussen,
27
28 S. G.; Choi, H.-J.; DeVree, B. T.; Sunahara, R. K. Structure and function of an irreversible agonist-
29
30 [bgr] 2 adrenoceptor complex. *Nature* **2011**, *469*, 236-240.
31
32
33 18. *Sybyl, X.* 1.0 Tripos Inc 1699 South Hanley Road. St Louis. Missouri 63144.
34
35 19. Barak, L. S.; Salahpour, A.; Zhang, X.; Masri, B.; Sotnikova, T. D.; Ramsey, A. J.; Violin, J.
36
37 D.; Lefkowitz, R. J.; Caron, M. G.; Gainetdinov, R. R. Pharmacological characterization of
38
39 membrane-expressed human trace amine-associated receptor 1 (TAAR1) by a bioluminescence
40
41 resonance energy transfer cAMP biosensor. *Mol. Pharmacol.* **2008**, *74*, 585-594.
42
43 20. Salahpour, A.; Espinoza, S.; Masri, B.; Lam, V.; Barak, L. S.; Gainetdinov, R. R. BRET
44
45 biosensors to study GPCR biology, pharmacology, and signal transduction. *Front. Endocrinol.*
46
47 (*Lausanne*) **2012**, *3*, 105.
48
49
50
51
52
53
54
55
56
57
58
59
60

TOC Graphic.

